

Corn with Vitazyme application—A Germination Study

Researcher: Steven David

Research organization: Sustainable Farming Solutions,
Perth, Western Australia

Location: Western Australia

Variety: Unknown


Experimental design: Corn seeds were treated with Vitazyme and compared with an untreated control treatment to determine the effect of this product on germination.

① Control ② Vitazyme

Vitazyme application: The treated seeds were treated on blotter paper with a 5% Vitazyme solution.

Growth results: Root and shoot growth were not measured, but can be clearly seen to be superior with the Vitazyme treatment. Note the accompanying photograph.

Conclusions: A corn seed treatment study with Vitazyme revealed that a 5% solution will dramatically enhance the development of roots and shoots.



Corn seeds treated with a 5% Vitazyme solution show much greater vitality on emergence than the untreated control. Note especially the improved rooting.



Corn with Vitazyme application—A Study on Synergism with WakeUp

Researchers: Steve Schmidt and Bert Schou, Ph.D.
Research organization: ACRES Research, Cedar Falls, Iowa
Location: Denver, Iowa
Variety: P0 589 AM
Planting date: May 18, 2018
Planting rate: 34,600 seeds/acre
Planting depth: 2 inches
Row spacing: 30 inches
Plot size (harvested area): 5x60 feet
Soil type: Dickinson fine sandy loam (2% organic matter, 6.3 pH, 9.1 meq/100 grams cation exchange capacity).
Experimental design: A small-plot corn trial, with four replications and six treatments, was prepared in east-central Iowa to evaluate the effect of Vitazyme, applied at different times, on corn yield. An additional treatment was added to evaluate the effect of WakeUp adjuvant as a synergist with Vitazyme.



A much more massive root system with the Vitazyme seed treatment produced an 8.0 bu/acre yield increase compared to the untreated control.


Treatment	Vitazyme			WakeUp	
	In-furrow oz/acre	V6 oz/acre	R1 oz/acre	Spring oz/acre	Summer oz/acre
1. Control	0	0	0	0	0
2. Vitazyme in-furrow	13	0	0	0	0
3. Vitazyme foliar/soil V6	0	13	0	0	0
4. Vitazyme foliar R1	0	0	13	0	0
5. Vitazyme foliar/soil V6 + R1	0	13	13	0	0
6. Vitazyme + WakeUp Spring in-furrow + Vitazyme + WakeUp Summer R1	13	0	13	4	4



Ear filling was enhanced considerably by a Vitazyme seed treatment at Denver, Iowa.

Fertilization: 90 lb/acre of N
Vitazyme application: (1) 13 oz/acre in-furrow (treatments 2 and 6); (2) 13 oz/acre to the leaves and soil at V6 on June 13 (Treatments 3 and 5); (3) 13 oz/acre to the leaves at R1 (silking; Treatments 4, 5, and 6).
WakeUp Spring application: 4 oz/acre in-furrow (Treatment 6).
WakeUp Summer application: 4 oz/acre at R1 (silking; Treatment 6)
Harvest date: October 17, 2018, using a MF 8 plot combine
Plant population results: The treatments ranged in population from 32,210 to 33,165 plants/acre, and did not differ significantly at P=0.05 (F-value = 0.8866 for replicates, and 0.6809 for treatments).


Grain moisture results:



**Decrease in grain moisture at harvest with Vitazyme:
0.5 to 1.3 percentage points**

Statistics	
LSD (P=0.05)	1.12%
Standard deviation	0.021%
CV.....	1.19
Replicate F	0.946
Treatment F	0.542

Grain yield results:



LSD (P=0.05)	14.1 bu/acre
Standard deviation ...	10.7 bu/acre
CV.....	5.12
Replicate F	0.350
Treatment F	0.590

Vitazyme in-furrow	+ 8.0 bu/acre (+4%)
Vitazyme at V6	+ 2.9 bu/acre (+1%)
Vitazyme at R1	- 1.8 bu/acre (-1%)
Vitazyme at V6 + R1	+ 8.5 bu/acre (+4%)
Vitazyme at IF + R1	+ 4.3 bu/acre (+2%)
+WakeUp IF + R1	

Conclusions: A small-plot research trial in eastern Iowa, using Vitazyme at different times, and together with WakeUp application in one treatment, revealed that corn yield responded to Vitazyme applied in-furrow (+ 4%), and also when applied at V6 and R1 (silking) at 13 oz/acre each time. The WakeUp applications with Vitazyme gave only a 2% yield increase, implying that there was little apparent synergism between the two products, though none of the yield values were significantly different. The R1 application alone gave a slightly negative, insignificant response. This treatment however, gave a slight reduction in grain moisture. These results corroborate the consistent effect of Vitazyme to increase corn grain yields in the United States Corn Belt.

Corn Vitazyme on Corn—A Synergism Study with WakeUp and Environoc 40IC



Researcher: Gilman Farley and Jerry Carlson

Research organization: Biodyne Midwest, Fort Wayne, Indiana, and Renewable Farming, LLC, Cedar Falls, Iowa

Location: eastern Iowa

Variety: unknown

Planting dates: unknown (2017 cropping year)

Experimental design: Eight field areas were treated in 2017 with different treatments, using WakeUp Spring, Vitazyme, and Environoc 40IC in combinations, to determine the yield of corn on an eastern Iowa farm. Two of the field areas were untreated controls. Samples of two to 10 acres were harvested from each of the eight areas and analyzed for yield using a combine monitor.

Product	Plot 1	Plot 2	Plot 3	Plot 4	Plot 5	Plot 6	Plot 7	Plot 8
	----- application in-furrow per acre -----							
1. WakeUp Spring	0	3 oz	0	3 oz				
2. Vitazyme	0	8 oz	0	8 oz				
3. Environoc 40IC	0	1 pint	0	1 pint	1 pint	0	0	0

Fertilization: according to soil test results, including chicken litter and gypsum

Vitazyme application: 8 oz/acre in-furrow to Plots 2, 3, 4, 5, 6, and 8.

WakeUp Spring application: This Renewable Farming product, a vegetable oil-based adjuvant, was applied at 3 oz/acre in-furrow to Plots 2, 3, 4, 5, 6, and 8.

Environoc 40IC application: This Biodyne product, containing various bacteria, nitrogen-fixing organisms, and


mycorrhizal fungi, was applied at 1 pint/acre in-furrow to Plots 2, 4, and 5.

Harvest date: unknown

Yield results: An 8-row combine with a calibrated yield monitor harvested areas of two to 10 acres of each treated field. The combine yield monitor maps were analyzed by Larry Eekhoff, a professional agronomist, and yield data was extracted from rows near each other to minimize soil effects. Plots having the same treatment are averaged in the table.

Treatment	Plot								Mean	Yield Change
	1	2	3	4	5	6	7	8		
	----- bu/acre -----									
1. Control	201.3	—	—	—	—	—	192.0	—	196.7	—
2. WakeUp Spring + Vitazyme	—	—	228.5	—	—	224.4	—	227.9	226.9	30.2(+15%)
3. WakeUp Spring + Vitazyme + Environoc 40IC	—	229.5	—	253.8	228.2	—	—	—	237.2	40.5(+21%)

Grain Yield



The plot yields for the control and WakeUp Spring + Vitazyme were very consistent, varying only 9.3 bu/acre and 4.1 bu/acre, respectively. Plot 4 of the three products combined yielded 25.6 bu/acre more than the lowest yield plot with that treatment, giving that treatment a 21% increase over the control and 6% more yield than the WakeUp + Vitazyme treatment.

Conclusions: This in-field study in eastern Iowa, comparing in-furrow applications at recommended rates of WakeUp Spring + Vitazyme and WakeUp Spring + Vitazyme + Environoc 40IC with the untreated control, revealed that these two combinations of products provided excellent synergisms for yield increases. These increases were 30.2 (15%) and 40.5 (21%) bu/acre for the respective treatments, revealing that an adjuvant, together with a biostimulant and appropriate microorganisms that fix nitrogen and help make nutrients available to plant roots, worked very well together. The surface tension reducing properties of WakeUp Spring to enhance cell membrane nutrient passage, plus the Environoc 40IC organisms which are triggered by the active agent of the Vitazyme, produced strong evidence of an excellent mutualism in this study. These yield increases were extremely profitable considering the low cost of the inputs, even for a year when grain prices were very low.



Corn with Vitazyme application—Mutualistic Effects with Environoc 401

Researcher: Gilman Farley

Research organization: Biodyne USA, Ft. Wayne, Indiana

Location: eastern Iowa

Variety: Unknown

Experimental design: A corn field was selected to evaluate the effectiveness of Vitazyme, together with Environoc 401, to improve grain yield as compared to an untreated area.


1 Control 2 Vitazyme + Environoc 401

Fertilization: 4 gal/acre of 6-24-6% N-P₂O₅-K₂O

Vitazyme application: 4 oz/acre (0.3 liter/ha)

Environoc 401 application: 16 oz/acre in-furrow. Environoc 401 is a selection of beneficial microbes that populate the rhizosphere for increased growth.

Yield results: A combine monitor was used for yield determinations. The moisture content of the corn grain varied within a narrow range; 16.4% to 17.1%. Two side-by-side comparisons were made in the field.




The Vitazyme + Environoc 401 effects on corn yield can be clearly noted on this combine monitor printout; see the darker green strip on the left half of the field.

Treatment	Yield bu/acre	Yield change bu/acre
Comparison 1:		
Control	228.8	—
Vitazyme + E 401	252.4	23.6 (+10%)
Comparison 2:		
Control	222.8	—
Vitazyme + E 401	235.9	13.1 (+6%)
Average:		
Control	225.8	—
Vitazyme + E 401	244.2	18.4 (+8%)

Overall increase in corn yield with Vitazyme: 8%

Conclusions: This on-farm corn yield study in north-central Iowa, using Vitazyme together with Environoc 401 in-furrow at planting, revealed that the yield was markedly improved, by 13.1 to 23.6 bu/acre, averaging 18.4 bu/acre (+8%). This result shows the excellent effectiveness of this program for corn farmers in Iowa.



Corn with Vitazyme application—An Evaluation on Effects with Mexican Corn Rootworm



Researcher: Lucino Frias, Eng.
Research organization: Quimica Lucava, S. A., Celeya, Guanajuato, Mexico
Farmer: Octavio Solorzano
Location: Estipac, Jalisco, Mexico
Variety: Pioneer
Planting date: June 15, 2018

Experimental design: A 1 hectare area of a corn field was treated with Vitazyme, and compared with the adjoining untreated control area to evaluate the effect of this product on the damage caused by the Mexican corn rootworm (*Diabrotica virgifera zea*) on plant growth and yield.

① Control ② Vitazyme

Fertilization: unknown

Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 6-leaf stage, 25 days after planting (July 10, 2018)

Growth and yield results:

Nine Days After Application (July 19)

Note more abundant healthier roots with Vitazyme.



The roots of the Vitazyme treated corn on the right reveal superior growth compared to the untreated control on the left.

Twenty Days After Application (July 30)

The stem diameter is greater, the height is greater, and the plants are more vigorous with Vitazyme.



Notice how much more growthy is the treated corn on the right, which displays more leaves, thicker stalks, and a darker green color due to more chlorophyll.

Fifty-Seven Days After Application (September 5)

There was damage present by Mexican corn rootworm (*Diabrotica virgifera zea*), which was causing much plant lodging. However, plants treated with Vitazyme, because of their greater root development, recovered better.



Rootworm-damaged corn treated with Vitazyme has more aggressively turned upwards to regain its stature, compared to the untreated control plant.

Eighty-Five Days After Application (October 3)

It was observed that, even with the Diabrotica (Mexican corn rootworm) and lodging issues, plants and ears from the Vitazyme-treated area had better filling, better size and better weight, than the untreated control.

The Day Before Harvest

At this time, three samples of representative ears were collected from each treatment and weighed as-is.

Vitazyme

Weight: 392 grams



401 grams



371 grams



Untreated Control

Weight: 268 grams




321 grams



314 grams




Corn Yield



Increase in yield with Vitazyme: 9%

Average Ear Weight



Increase in ear weight with Vitazyme: 29%

Conclusions:

- Vitazyme treated plants showed healthier and more abundant roots, thicker stems, and taller and more vigorous plants than the untreated control.
- Vitazyme treated plants rebounded stronger from lodging caused by the attack of Mexican corn rootworm (*Diabrotica virgifera zea*), in comparison with the untreated control.
- Plants and ears from the area treated with Vitazyme had better filling, better size, and 29% greater ear weight than those of the untreated control.
- Vitazyme caused a 9% yield increase (0.63 tonne/ha) with only a single 1 liter/ha foliar spray.

Income Results: At \$0.20/kg of grain, Vitazyme produced \$123/ha greater income.



Corn with Vitazyme application

Researcher: Gustavo Figueroa Ramirez
Location: Cofradia de Lepe, Municipality of Atengo, El Fresno Ranch, Jalisco, Mexico

Variety: Unknown

Planting date: June 28, 1018

Experimental design: A 4-hectare portion of a corn field was treated with two different Vitazyme application regimes to evaluate the effect of the product on corn growth and yield in a field setting.

1 Control 2 Vitazyme

Fertilization: unknown

Vitazyme application: (1) Seed treatment on June 28 of 0.25 liter of Vitazyme in 0.25 liter of water (50% solution) to coat seeds for 1 hectare; (2) 1 liter/ha sprayed by backpack sprayer at the 6-leaf stage; (3) 1 liter/ha sprayed by backpack sprayer at the 9-leaf stage. One small area received only treatment (3).

Growth observations: It was noted that the Vitazyme treated plants were much taller and had more leaf area than the control plants, with larger ears and considerably greater root development.



The Vitazyme treated corn displays excellent ear development, with filling to the tip and large kernels, giving a 35% greater yield than the control.

Plant height and leaf growth were enhanced greatly in this Mexican corn trial.



Preliminary Yield Results			
	Control	Vitazyme	Change
Complete plant weight, kg	1.02	2.72	+1.70 (+167%)
Ear weight with husk, grams	340	530	+190 (+56%)
Ear weight without husk, grams	250	350	+100 (+40%)

Conclusions: This Mexican corn study, using a seed treatment and two foliar treatments of Vitazyme, showed remarkable responses in terms of plant size and ear weight. Harvest is expected in January of 2019, at which time yield data will be available.



Corn with Vitazyme application

Researcher: James Anderson
Research organization: J & H Distributing, Belgrade, Minnesota
Location: Pederson Farms, Spicer, Minnesota
Variety: P9929AMXT and DCK 45-65
Planting date: unknown
Soil type: silty clay loam
Experimental design: A corn field of 132.2 acres was treated in part (89.3 acres) with Vitazyme in-furrow, to determine the effect of the product on corn yield.
Fertilization: unknown

1 Control 2 Vitazyme

Vitazyme application: 13 oz/acre (1 liter/ha) in-furrow at planting
Crop season weather: a cold and late spring, with a summer of average temperatures and above-average rainfall

Treatment	Yield bu/acre	Yield change bu/acre
1. Control	200.9	—
2. Vitazyme	208.6	7.7 (+4%)




The Vitazyme seed treated plants are larger with considerably better developed root systems.



More full kernel development of the Vitazyme treated plants resulted in a higher yield of 8 bu/acre, a highly profitable application despite low corn prices.

Yield results: Both the control and treated grain were harvested at 22% moisture. A combine monitor was used to record yields.



Conclusions: An on-farm corn trial in central Minnesota proved that Vitazyme, applied in-furrow at 13 oz/acre (1 liter/ha) increased the grain yield by 7.7 bu/acre (4%), showing the high profitability of this amendment for corn farmers in the northern Corn Belt.

Increase in corn yield with Vitazyme: 4%



Corn with Vitazyme application

Researcher: David Clay Ph. D., and Graig Reicks

Research institution: Department of Plant Sciences, South Dakota State University, Brookings, South Dakota

Location: South Dakota State University, Field Research Facility, Aurora, South Dakota

Variety: DKC 47-47

Planting date: May 15, 2018

Planting rate: 30,750 seeds/acre

Planting depth: 2 inches

Row spacing: 30 inches

Soil type: Brandt silty clay loam

Experimental design: A replicated, randomized complete block design, with four replications, was set up in a small-plot corn trial, plots being six rows wide (15 feet) and 20 feet long. At harvest, 20 feet of the two middle rows were machine-harvested with a plot combine. Three nitrogen rates and three different times of application were utilized. Grain yield was measured, and nitrogen efficiency and water efficiency were to be determined through ¹⁵N and ¹³C analyses.

Vitazyme application	Nitrogen rate, lb/acre		
	0	75	125
	Treatment number		
Control	1	2	3
Seed only	4	5	6
Foliar V6 only	7	8	9
Seed + Foliar V6	10	11	12
Seed + Foliar R1	13	14	15

Fertilization: Nitrogen fertilizer was applied by hand as urea (46%N) to appropriate plots on June 1, when the corn was in the V1/V2 (one to two-leaf) stage.

Vitazyme application: Seed treatment (at planting on May 15): 156 ml of Vitazyme per 12.3 lb of seed, or 8 oz/acre sprayed on the seeds and mixed by hand in totes. **Foliar treatment** (on June 22 at V6, and on July 27 at R1): 13 oz/acre using a 15 gal/acre spray solution with a non-ionic surfactant (0.25%) and a backpack sprayer.

Harvest date: unknown



Vitazyme on the seeds, as well as on the leaves, produced better filled ears with larger kernels, as can be seen in this 2018 nitrogen efficiency study.




Notice the excellent root development with Vitazyme application at 75 lb/acre of nitrogen. This has led to better ear filling, as seen in the other photo.

Yield results:

Nitrogen Fertilizer Effect			
Vitazyme	N rate	Grain yield ¹	Yield change ²
	lb/acre	bu/acre	bu/acre
1. None	0	153 e	—
2. None	75	197 a	+ 44 (+29%)
3. None	125	195 ab	+ 42 (+27%)
4. Seed	0	170 cde	—
5. Seed	75	194 ab	+ 24 (+14%)
6. Seed	125	200 a	+ 30 (+18%)
7. Foliar V6	0	171 bcde	—
8. Foliar V6	75	197 a	+ 26 (+15%)
9. Foliar V6	125	190 abcd	+ 19 (+11%)
10. Seed + Foliar V6	0	167 de	—
11. Seed + Foliar V6	75	199 a	+ 32 (+19%)
12. Seed + Foliar V6	125	197 a	+ 30 (+18%)
13. Seed + Foliar R1	0	—	—
14. Seed + Foliar R1	75	198 a	n.a. ³
15. Seed + Foliar R1	125	193 abc	n.a. ³

¹Means followed by the same letter are not significantly different at P=0.05. ²Change vs. the O-N rate for that Vitazyme treatment. ³Data is not available.

Average Fertilizer Effect¹




¹Averaged over all Vitazyme treatments.

Increase in yield with 75 lb/acre N:
31.7 bu/acre (+19%)

Increase in yield with 125 lb/acre N:
29.7 bu/acre (+18%)

Vitazyme Effects at 0 N



Vitazyme Treatment	Grain yield bu/acre	Yield change bu/acre
None	153	—
Seed	170	17 (+11%)
Foliar V6	171	18 (+12%)
Seed + Foliar V6	167	14 (+9%)

Increase in yield with seed treatment: 11%

Increase in yield with foliar treatment at V6: 12%

Increase in yield with seed + foliar treatment at V6: 9%

Vitazyme effects at the 75 and 125 lb/acre N rates were not significant, likely because the corn crop had reached its maximum yield potential at about 195 to 200 bu/acre.

Conclusions: This South Dakota State University corn study, using three nitrogen fertilizer rates, plus four Vitazyme application regimes, revealed that, with no added nitrogen, Vitazyme elevated the grain yield by 9 to 12% above the untreated control, by amounts of 14 to 18 bu/acre. At nitrogen rates of 75 and 125 lb/acre, however, the yields of all treatments were similar, varying from 190 to 200 bu/acre, so Vitazyme effects were not detected because the maximum yield potential had been reached even at 75 lb of N/acre. This medium application rate outyielded the high nitrogen rate by 2 bu/acre. It would have been proper during this trial to have utilized lower fertilizer nitrogen rates, such as 25 and 50 lb/acre, to evaluate fertilizer efficiency effects of Vitazyme at these lower rates.

Evaluations of nitrogen and water efficiency will be conducted on these 15 treatments during the coming months, and will be reported in upcoming data summaries.



Corn with Vitazyme application

Researcher: E. Bruce Kirksey, Ph. D.
Research organization: Agricenter International, Memphis, Tennessee
Location: Memphis, Tennessee
Variety: DK 65-19
Soil type: Falaya clay loam; good fertility and drainage; pH=6.4
Row spacing: 30 in
Plant population: 32,000 seeds/acre
Planting date: May 14, 2018
Experimental design: A small-plot replicated field corn trial was prepared using two Vitazyme applications, to determine the yield and profitability of this program compared to the untreated control. Four replications of plots 10 x 30 ft were used.


1 Control 2 Vitazyme

Fertilization: unknown
Vitazyme application: (1) 13 oz/acre (1 liter/ha) in-furrow at planting; (2) 13 oz/acre (1 liter/ha) sprayed on the leaves and soil at V6

Yield results: The corn grain from each plot was shelled and weighed on September 14, using an Almaco plot combine. Two rows 25 ft long were harvested.

Treatment	Yield ¹ bu/acre	Yield change bu/acre
Control	135.5 b	—
Vitazyme	173.7 a	38.2 (+28%)
Standard error	4.1	

¹Means followed by the same letter are not significantly different at P=0.05.



Increase in corn grain with Vitazyme: 38.2 bu/acre (+28%)

Income results:

Income results	
Added income from the two applications...	\$143/acre
Added costs	\$17/acre
Net profit	\$126/acre
Return on investment	7.41:1

Conclusion: A small-plot replicated corn trial in Tennessee revealed a dramatic increase in shelled corn yield with in-furrow and foliar(V6) Vitazyme applications. The yield increase was 38.2 bu/acre above the untreated control treatment, or 28% more. This increase netted \$128/acre more income, or a return on investment of 7.41:1. This increase, if extrapolated to a 1,000-acre corn farming operation, would give the farmer \$126,000 more profit, from product costing him only about \$17,000.



Corn with Vitazyme application

Researcher: V. V. Plotnikov

Research organization: Plant Designs, Inc., Rochester, New York, and Agro Expert International, Kaharlyk, Ukraine

Location: Drabiv District, Cherkasy Region, Novomykolaivka Village, Private Enterprise Opal, Ukraine

Variety: ES Sensor, FAO 370

Planting date: April 30, 2018

Previous crop: winter wheat

Soil type: typical chernozem (humus = 4.1%)

Planting rate: 70,000 seeds/ha

Field preparation: disking to 6-8 cm, plowing to 22-24 cm, cultivation to 5-6 cm

Experimental design: A corn field was treated in part with two applications of Vitazyme, and compared with the untreated portion of the field, to evaluate the effect of the product on the yield and profitability of corn grain.

① Control ② Vitazyme

Fertilization: 123-30 kg/ha of N-P₂O₅ at planting; 10-26-26 kg/ha of N-P₂O₅-K₂O at planting

Vitazyme application: (1) 1 liter/ha sprayed on the soil pre-plant on April 27; (2) 0.5 liter/ha sprayed on the leaves and soil at the 7 to 8-leaf stage on May 30

Yield results:


Treatment	Yield tonnes/ha	Yield change tonnes/ha
Control	11.2	—
Vitazyme	12.8	1.6 (+14%)

¹Means followed by the same letter are not significantly different at P=0.05.

Increase in grain yield with Vitazyme: 14%



Corn produced using Vitazyme shows better filling and ear size as a result of superior nutrient uptake by an enlarged root system and activated rhizosphere.



Income results: An extra yield of 1.6 tonnes/ha produced an income increase of \$281/ha.

Conclusion: This Ukrainian corn trial, using 1 liter/ha and 0.5 liter/ha sprays pre-plant and at the 7 to 8-leaf stage, respectively, resulted in an excellent 14% yield increase, which brought \$281/ha more income. This program is highly effective for corn producers in Ukraine.



Corn with Vitazyme application—A Study on Synergism with WakeUp Summer

Researchers: Steve Schmidt and Bert Schou, Ph.D.

Research organization: ACRES Research, Cedar Falls, Iowa

Location: Denver, Iowa

Variety: Wyffels W5440

Planting date: May 16, 2017

Planting rate: 35,400 seeds/acre

Planting depth: 2.25 inches

Row spacing: 30 inches

Soil type: Organic matter, 3.0%; pH, 7.3; cation exchange capacity, 11.03 meg/100g.

Field preparation: strip tillage on April 24, 2017

Previous crop: soybeans

Experimental design: A medium-sized randomized complete block design with five replicates was established on a farmer's field, total plot size being 15 x 60 feet and the harvested area of each plot being 5 x 60 feet (0.00689 acre). Vitazyme and an adjuvant (WakeUp Summer) were applied alone and together to determine the effect of each treatment on the grain yield.

- 1 Control
- 2 Vitazyme
- 3 WakeUp Summer
- 4 Vitazyme + WakeUp Summer

Fertilization: Starter mix (4-Star Mix) at planting: 5 gal/acre of MPK (0-52-34% N-P₂O₅-K₂O), 3 gal/acre of K-row (0-0-23-8 % N-P₂O₅-K₂O-S), 1 quart/acre of humate, and 4 oz/acre of Supertrace.

Vitazyme application: (1) 13 oz/acre (1 liter/ha) in-furrow at planting (May 16) and (2) 13 oz/acre (1 liter/ha) on the leaves and soil at R1 on June 26.

WakeUp Summer application: (1) 4 oz/acre in-furrow at planting (May 16) and (2) 4 oz/acre on the leaves and soil at R1, on June 26.

Herbicides: Pre-emergence (applied May 16), 5.6 oz/acre Corvus, 1 quart/acre Roundup, 1 lb/acre Atrazine, 1 pint/acre 2, 4-D, 1 lb/acre AMS, in 15 gal/acre of water. Post-em organs (applied June 16), 1 oz/acre Impact, 0.25 lb/acre Atrazine, 8 oz/acre MSO, and 2 lb/acre AMS.

WakeUp Summer: a vegetable-based adjuvant

Growing season weather: favorable throughout the growing season



The Vitazyme treated corn showed the greatest rooting and overall growth in this Iowa study, from two applications, at planting and at V8.

Population results: Population counts were made on September 14. All treatments varied in population from 31,376 to 32,509 plants/acre, and were not significantly different.

Grain moisture results: Grain moisture varied from 19 to 20%, and were not significantly different.


Grain yield results: The plots were harvested on October 28, 2017 using a plot combine, during which grain weight and moisture content were recorded.



Note the larger, better filled ears from the Vitazyme treated corn in this study on the right, versus the control corn on the left.

Treatment	Yield ¹ bu/acre	Yield change bu/acre
1. Control	227 b	—
2. Vitazyme	242 a	15 (+7%)
3. WakeUp Summer	240 a	13 (+6%)
4. Vita + WakeUp	238 ab	11 (+5%)

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test. LSD 0.05=12.5 bu/acre.



Conclusions: This small-plot study on corn near Denver Iowa, revealed that both Vitazyme and WakeUp Summer significantly improved corn grain yield, by 6% for WakeUp and 7% for Vitazyme. The combination of the two products also increased grain yield -by 5%- but this increase was slightly less than for each product by itself. Both Vitazyme and WakeUp Summer are shown to significantly improve corn yield when applied at 13 oz/acre (1 liter/ha) in furrow and at R1, and at these same growth stages for WakeUp, but at the 4 oz/acre rate.



Corn with Vitazyme application—A Nitrogen Efficiency Study

Researcher: Dr. David Clay and Graig Reicks
Research institution: Department of Plant Sciences, South Dakota State University, Brookings, South Dakota
Location: South Dakota State University, Field Research Facility, Aurora, South Dakota
Variety: DKC 49-72 (99-day hybrid)
Planting date: May 14, 2016
Planting rate: 32,000 seeds/acre
Planting depth: 2 inches
Row spacing: 30 inches
Soil type: Brandt silty clay loam
Experimental design: A replicated, randomized complete block design, with three replications, was established in a small-plot corn trial. Plots were six rows wide (15 feet) and 20 feet long, and at harvest 20 feet of the two middle rows were hand-harvested. Three nitrogen rates were utilized, along with Vitazyme applied on the seeds or foliar, to determine the effect of this product on corn yield, protein (nitrogen) levels, and nitrogen and water efficiency.



The research plots at South Dakota State University produced excellent yield and nitrogen efficiency data for 2016.



Notice the much improved root development when Vitazyme is applied to the seeds and leaves.

Nitrogen rate lb/acre	Vitazyme application			
	None	Seeds	Leaves	Seeds + Leaves
0	1	4	7	10
75	2	5	8	11
125	3	6	9	12

Fertilization: Nitrogen fertilizer was applied by hand as urea (46%) to appropriate plots on June 13, 2016, when the corn was at the V4 (four-leaf) stage.
Vitazyme application: Seed treatment (at planting on May 14): 112 ml of Vitazyme per 10 lb of seed, or 7.4 oz/acre sprayed on the seeds and mixed on a tarp; **Foliar treatment** (on June 29 at V 8-9) at 13 oz/acre, using a 15 gal/acre spray solution, with a non-ionic surfactant (0.25% v/v) and a backpack sprayer.

¹⁵N and ¹³C determinations: A mass spectrometer was used at the Department of Plant Science, South Dakota State University.

Harvest date: hand harvested on October 31, 2016


Yield, nitrogen efficiency, and water efficiency results: Ears were dried at 140° F and weighed daily until moisture loss ceased. Grain was then removed by a portable sheller and weighed. Yields were adjusted to 15% moisture. Yield loss due to water and nitrogen stress were then calculated using ¹⁴N, and ¹⁵N, and ¹³C formulas.

Vitazyme Seed Treatment Effect on Corn Yield and Nitrogen Grain Use Efficiency

Treatment	Nitrogen fertilizer bu/acre	Corn yield ¹ bu/acre	Corn grain use efficiency ² %
1. None	125	220 a	43.3 b
	75	197 b	40.3 b
	0	181 bc	—
2. Vitazyme on seeds	125	218 a	42.6 b
	75	218 a	58.0 a
	0	162 c	—

¹Means followed by the same letter are not significantly at P=0.05. ²Means followed by the same letter are not significantly different at P=0.10. The calculations used to determine this value are based upon 14N and 15N ratios in the grain, and will not be presented in this report.


Vitazyme Seed Treatment Effect on Corn Yield



Yield increase with Vitazyme at 75lb/acre of nitrogen: 21 bu/acre

Corn yield with 75 lb/acre of N, using a 40% reduction in N, yielded the same as for the 100% (125 lb/acre) N rate.

Vitazyme Seed Treatment Effect on Corn Grain Use Efficiency at 75 lb/acre Nitrogen




Vitazyme Seed Treatment Effect on Corn Yield and Water and Nitrogen Stress


Vitazyme seed treatment	Nitrogen fertilizer rate lb/acre	Grain yield lb/acre	Yield loss water stress lb/acre	Yield loss nitrogen stress lb/acre	Grain nitrogen %
Yes	125	218 a*	8.7 b*	9.2 b*	1.31 a*
Yes	75	218 a	8.7 b	8.8 b	1.30 a
Yes	0	162 b	40.5 a	31.8 a	1.09 b
No	125	220 a*	10.3 c**	4.1 b**	1.47 a*
No	75	197 b	20.7 b	16.9 a	1.29 b
No	0	181 b	32.2 a	21.6 a	1.14 c

*Means followed by the same letter are not significantly different at P=0.05. **Means followed by the same letter are not significantly different at P=0.10.

Water Stress Yield Loss (WSYL)



Nitrogen Stress Yield Loss (NSYL)



<i>Reduction in Water Stress Yield Loss</i>		
	<i>Vitazyme</i>	<i>No Vitazyme</i>
<i>At 75 lb/acre N</i>	<i>79%</i>	<i>26%</i>
<i>At 125 lb/acre N</i>	<i>79%</i>	<i>68%</i>

<i>Reduction in Nitrogen Stress Yield Loss</i>		
	<i>Vitazyme</i>	<i>No Vitazyme</i>
<i>At 75 lb/acre N</i>	<i>72%</i>	<i>22%</i>
<i>At 125 lb/acre N</i>	<i>71%</i>	<i>81%</i>

Income results: At \$4.00/bu of corn grain, the extra 23 bu/acre at 75 lb/acre of N is \$92.00. A savings of 50 lb/acre N, at a cost of \$0.27/lb of urea, is \$13.50. That \$105.50 greater income, less an approximate cost at \$7.00/acre for Vitazyme, equals a Benefit: Cost of 15.1.

Benefit Cost of 15.1

Conclusions: A replicated small plot corn study conducted by South Dakota State University in 2016, near Aurora, South Dakota, using ¹⁵N and ¹³C analytical methods, revealed that Vitazyme on the seeds greatly improved the efficiency of both

water and nitrogen use. This was shown in particular by the changes in efficiency at 75 lb/acre (40% N reduction) of nitrogen application: treated seeds reduced water stress yield loss by 79%, while nitrogen stress yield loss was reduced by 72%. Grain use efficiency (the percent of applied fertilizer actually going into the grain) at 75 lb/acre of N fertilizer for the Vitazyme treated seed was 58%, while the untreated seed efficiency was 40.3%. These effects of the Vitazyme seed treatment translated into virtually the same yield for Vitazyme on the seeds at 75 lb/acre of N (218 bu/acre) as for

untreated seeds at 125 lb/acre of N (220 bu/acre). At 75 lb/acre of N, the treated seeds yielded 23 bu/acre more than the untreated seeds. Even with depressed corn prices of around \$4.00/bu the benefit: cost ratio of the Vitazyme seed treatment with reduced N application would be 15.1:1. Grain nitrogen was also significantly increased by Vitazyme on the seeds.

The Vitazyme foliar treatment did not yield significantly positive results in this trial. Reasons for this lack of foliar response are not known, since a trial in 2014 did produce a significant yield increase of 9 bu/acre.



Corn and Soybeans with Vitazyme application

Research organization: University of Missouri, School of Natural Resources and Division of Plant Sciences, Columbia, Missouri, USA

Researchers: Robert J. Kremer, Manjula V. Nathan, Paul W. Syltie, Tim Reinbott, Kelly Nelson, and Xiaowei Pan

Introduction: Biostimulants enhance nutrient uptake and crop quality; however, knowledge of effects on soil health is limited (Kremer 2017a). The biostimulant Vitazyme provides several modes of action mediated by multiple active brassinosteroids. Attributes of brassinosteroid-based biostimulants include enhanced root development, improved nutrient uptake efficiency, disease suppression, and environmental stress tolerance (Hayat & Ahmad 2011). Limited research has shown that brassinosteroids may regulate pesticide metabolism in plants (Zhou et al. 2015); however, no information is available on the potential for overcoming stress of the herbicide glyphosate on rhizosphere biology in transgenic (genetically-modified, [GM]) cropping systems. Glyphosate affects rhizosphere microbial diversity and activity, increased fungal root conlonization, decreased beneficial bacterial components, and detrimental effects on crop root growth (Kremer & Means 2009; Wagner et al. 2003). Vitazyme was applied in soybean and maize field trials in Missouri USA during 2014-2017, with and without glyphosate application.





Objectives:


- Determine effects of Vitazyme on selected soil health indicators
- Determine effects of Vitazyme on rhizosphere biology in transgenic soybean and maize receiving glyphosate in field trials
- Determine potential effects of Vitazyme on crop growth in soils with residual glyphosate contents

Experimental design:

- Field trials conducted at Columbia (2014) and Novelty (2016-17), Missouri, on Mexico sil soil (fine, smectitic, sesic Vertic Epiaqualfs), maize and soybean planted conventionally using minimum tillage; plants and soils collected at R2 soybean and V10 maize growth stage
- Root-colonizing *Fusarium* assessed by selective culture technique (Levesque et al. 1993)
- Rhizosphere pseudomonads determined using S1 agar medium (Gould et al. 1985)
- Indoleacetic acid-producing (IAA) bacteria detected on nitrocellulose membranes reacted with Salkowski reagent for color development (Bric et al. 1991)
- Mn-transforming bacteria detected on Gerretsen's medium (Huber & Graham 1992)
- Glucosidase activity detected using enzyme assay of Eivazi & Tabatabai (1988)
- Soil microbial components and biomass determined using phospholipid fatty acid (PLFA) analysis (Buyer & Sasser 2012; Pritchett et al. 2011)


Soybean Results:





Note how indole acetic acid bacteria, which are detected in the petri dishes by the pink halos they form around the bacteria, are much more numerous in the Vitazyme treated soybean soil inoculum. The bacteria are very beneficial to plant growth.

Soybean Yield Projection With Soil Glyphosate




Corn Results:




Note the highly beneficial effect of Vitazyme in enhancing root growth and ear development in this University of Missouri trial.


Rhizosphere Pseudomonads




Maize Root Dry Mass



IAA-producing Bacteria




Fusarium




With corn rhizosphere soil samples, the prevalence of Fusarium fungi is much higher when glyphosate is applied to the leaves, but when Vitazyme is added with the glyphosate these fungi are held in check. The same occurs with soybeans.

Mn Transforming Bacteria



PLFA Nmole 2016



Overall results:

Vitazyme-improved soil health indicator and overcame glyphosate effects on maize and soybean at all sites across years:

- Root growth (dry biomass) improved while colonization by potentially pathogenic *Fusarium* fungi was significantly ($p < 0.05$) reduced, root nodulation on soybean improved or did not decrease with glyphosate + Vitazyme.
- Rhizosphere fluorescent pseudomonads, which contribute to beneficial plant growth-promoting functions, and IAA-producing bacteria, significantly increased in both Vitazyme-treated crops.
- Maintained high Mn reducers to Mn oxidizers ratio, notably in presence of glyphosate, suggesting adequate available Mn (reduced) for plant and microbial uptake and metabolism, and limiting unavailable, immobilized Mn (oxidized) associated with glyphosate and oxidizing bacteria
- Maintained soil microbial community composition (PLFA groups) and total microbial biomass in glyphosate treated crops at similar levels of control, non-glyphosate soils, which tended to improve mycorrhizal component
- Enhanced glucosidase activity, indicator of soil microbial function, in both control and glyphosate treatments

Glyphosate Soil Residues and Grain Yields:

- Residual glyphosate contents in soil ranged from 0 to >1000 ug/kg (ppb) of soil, which was randomly distributed among plots at field sites, regardless of glyphosate application timing
- Soybean grain yields increased 1 to 4% (data not shown)
- No relationship between yields of individual plots with soil glyphosate content was found for maize (2016)
- Apparent relationship between soybean yield in individual plots with soil glyphosate-tendency for lower yield with higher soil glyphosate; Vitazyme may offset potential yield decline due to glyphosate buildup in Mexico silt loam (6 of 8 plots)

Conclusions:

- A polyphasic microbial analysis multiple assessments of sensitive soil health indicators, previously used to evaluate GM crop effects on soil biology and ecology (Kremer & Means 2009), was successfully adapted for evaluating the biostimulant Vitazyme as a means of suppressing effects of glyphosate on root growth and rhizosphere biology in (GM) cropping systems and maintaining or improving soil health.
- Soil microbial diversity based on PLFA analyses was restored by Vitazyme in soils planted to maize and soybean treated with glyphosate; a high microbial diversity is essential to maintain a stable ecosystem and crop productivity (Grayston et al. 1998).
- Glyphosate residue buildup in soils under GM crops is a reality (Kremer 2017b) and may affect crop growth and yields in future seasons. Biostimulants such as Vitazyme might offset effects of residual glyphosate, however more evaluation under rigorous experimental conditions is required to confirm effect.
- Results from this project demonstrate that biostimulants can be a major management factor for addressing recurrent productivity problems and declining soil health associated with GM crops in current crop production systems.

References:

- Bric, J.M., Bostock, R.M., Silverstone, S.E., 1991.** Rapid in situ assay for indoleacetic acid production by bacteria immobilized on a nitrocellulose membrane. *Applied & Environmental Microbiology* 57:535-538.
- Buyer, J.S., Sasser, M., 2012.** High throughput phospholipid fatty acid analysis of soils. *Applied Soil Ecology* 61:127-130.
- Eivazi, F., Tabatabai, M.A., 1988.** Glucosidases and galactosidases in soils. *Soil Biology & Biochemistry* 20:601-606.
- Gould, W.D., Hagedorn, C., Bardinelli, T.R., Zablutowicz, R.M., 1985.** New selective medium for enumeration and recovery of fluorescent pseudomonads from various habitats. *Applied & Environmental Microbiology* 49:29-32.
- Grayston, S.J., Wang, S.Q., Campell, C.D., Edwards, A.C., 1998.** Selective influence of plant species on microbial diversity in the rhizosphere. *Soil Biology & Biochemistry* 30:369-378.
- Hayat, S., Ahmad, A. (eds.), 2011.** *Brassinosteroids: A Class of Plant Hormone*. Springer, London.
- Huber, D.M., Graham, R.D., 1992.** Techniques for studying nutrient-disease interactions. In Singleton, L.L. et al. (eds.), *Methods for Research on Soilborne Phytopathogens*, APS Press, St. Paul, MN. pp. 204-214.
- Kremer, R.J., 2017a.** Biotechnology impacts on soil and environmental services. In Al-Kasai, M., Lowery, B. (eds.), *Soil Health and Intensification of Agroecosystems*. Elsevier, New York. pp. 353-375.
- Kremer, R.J., 2017b.** Soil and environmental health after twenty years of intensive use of glyphosate. *Advances in Plant & Agricultural Research*. 6(5):000224, doi: 10.15406/apar2017.0600224.
- Kremer, R.J., Means, N.E., 2009.** Glyphosate and glyphosate-resistant crop interactions with rhizosphere microorganisms. *European Journal of Agronomy* 31:153-161.
- Levesque, C.A., Rahe, J.E., Eaves, D.M., 1993.** Fungal colonization of glyphosate-treated seedlings using a new root-planting technique. *Mycological Research* 97:299-306.
- Pritchett, K.A., Kennedy, A.C., Cogger, C.G., 2011.** Management effects on soil quality in organic vegetable

systems in western Washington. Soil Science Society of America Journal 75:605-615.

Wagner, R., Kogan, M., Prada, A.M., 2003. Phytotoxic activity of root absorbed glyphosate in corn seedlings (*Zea mays* L.). Weed Biology and Management 3:228-232.

Zhou, U., Xia, X., Yu, G., et al., 2015. Brassinosteroids play a critical role in the regulation of pesticide metabolism in crop plants. Scientific Reports 5:9018, doi:10.1038/srep09018



Corn with Vitazyme application

Researcher: Vadim Plotnikov
Research organization: PE "Bondarenko", Ukraine, Plant Designs, New York, USA, and Agro Expert International, Ukraine
Location: Rozdilna District, Odessa Region, Rozdilna Village, Ukraine
Variety: KVS Kaifus, FAO 330
Seeding rate: 60,000 seeds/ha
Planting date: May 4, 2017
Previous crop: wheat
Soil type: typical Chernozem; humus = 3.9%
Soil preparation: disking to 6-8 cm, plowing to 22-24 cm, harrowing to 5-6 cm
Experimental design: A corn field was divided into Vitazyme treated and untreated control areas to determine the efficacy of this product in promoting yield increases.

1 Control 2 Vitazyme


Fertilization: 50-70 kg/ha of N-S broadcast before planting
Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 8 to 9-leaf stage
Growing season weather: dry
Yield results:



Note the improved ear development with Vitazyme treatment in Ukraine.

Vadim Plotnikov examines corn sampled from a Vitazyme treatment in Ukraine. Such increases have been highly profitable in the country.

Treatment	Grain yield tons/ha	Yield change ton/ha
1. Control	5.6	—
2. Vitazyme	7.1	1.5 (+27%)



Increase in grain yield with Vitazyme: 27%

Income results: At a price of \$164.67/ton of corn grain, the added 1.5 tons/ha gave an additional \$247/ha income.

Conclusions: A corn trial in southern Ukraine, utilizing 1 liter/ha sprayed on the leaves and soil at the 8 to 9-leaf stage, revealed that the grain yield responded excellently by increasing 1.5 tons/ha (27%). This increase resulted in a profit enhancement of \$247/ha, showing how this program is an excellent addition to corn farmers' cultural programs in Ukraine.





Corn with Vitazyme application

Researcher: Vadim Plotnikov
Research organization: PJSC "Zlagoda", Ukraine, Plant Designs, New York, USA, and Agro Expert International, Ukraine
Location: Onufriivka District, Kirovograd Region, Vyshnivtsi Village, Ukraine
Variety: Cobalt, FAO 320
Seeding rate: 70,000 seeds/ha
Planting date: May 3, 2017
Previous crop: wheat
Soil type: dark-brown podzolic; humus = 2.6%
Soil preparation: disking to 6-8 cm, plowing to 22-24 cm, cultivation to 5-6 cm
Experimental design: A corn field was divided into Vitazyme treated and control portions to discover the efficacy of this product in promoting grain yield.

1 Control 2 Vitazyme

Fertilization: 115 kg/ha of N broadcast before planting, and 10-26-26 kg/ha N-P₂O₅-K₂O in-furrow at planting


Vitazyme application: 1 liter/ha sprayed on the leaves and soil on June 5, 2017, at the 7 to 8-leaf stage
Growing season weather: dry
Yield results:

Treatment	Grain yield tons/ha	Yield change ton/ha
1. Control	8.0	—
2. Vitazyme	9.7	1.7 (+21%)

Yield increase with Vitazyme: 21%

Conclusions: A Ukrainian large-scale field trial with corn in 2017, using a single 1 liter/ha foliar/soil application in early June, revealed that grain yield was increased by 1.7 tons/ha (21%), which gave an additional income of \$283/ha. These results show the considerable efficacy of utilizing this program for more profitable corn culture in Ukraine.

Income results: At \$166.47/ton, the extra 1.7 tons/ha yielded \$283/ha greater income.





Corn with Vitazyme application—A Soil Microbiology Study with Roundup (Glyphosate)



Dr. Nathan displays the Vitazyme treated corn ears on the right, and the control ears on the left. The difference in kernel development is obvious.



Note the superior rooting and ear development for the Vitazyme treatment at the Greenley Center trial.

Researcher: Manjula Nathan, Ph.D., and Robert Kremer, Ph.D.
Research institution: University of Missouri Department of Plant and Soil Sciences, and the USDA-ARS, Columbia, Missouri
Location: Greenley Memorial Research Center, Novelty, Missouri
Variety: Unknown
Soil type: clayey
Row spacing: 30 inches
Experimental design: A small plot corn trial was conducted using four replications and four treatments, to evaluate the effect of Vitazyme and Roundup (glyphosate), alone and in combination, on the yield of corn, as well as the effects of these products on the rhizosphere soil microbial populations of Fusarium and beneficial bacteria. Soil residues of glyphosate were also evaluated.

Treatment	In-furrow bu/acre	Foliar, 8-leaf stage bu/acre
1. Control	0	0
2. Glyphosate	0	36 oz/acre
3. Vitazyme	13/oz/acre	13 oz/acre
4. Vitazyme + Glyphosate	13/oz/acre (Vitazyme only)	13 oz/acre + 36 oz/acre



Fertilization: Nitrogen, phosphorus, and potassium were applied before planting according to soil test recommendations.
Vitazyme application: See the treatment plan.
Glyphosate application: See the treatment plan.
Weed control for the control and Vitazyme treatments: These plots were sprayed with herbicides before planting.
Growing season weather: favorable for corn growth
Yield results:

Treatment	Yield bu/acre	Yield change bu/acre
1. Control	229	—
2. Glyphosate	237	8 (+3%)
3. Vitazyme	238	9 (+4%)
4. Vitazyme + Glyphosate	239	10 (+4%)
P>F	0,079	

Increase in corn yield




Glyphosate3%
Vitazyme4%
Vitazyme + Glyphosate ...4%

Yield differences were almost significant at P=0.05, with the combined products yielding the most.



For the Vitazyme and glyphosate yield analyses, all plots treated with that material were compared with all plots that were not treated with it.

Soil microbial results:



Changes in Fusarium Colonization

Glyphosate +20%

Vitazyme -4%

Vitazyme + Glyphosate... -11%

Changes in Fluorescent Pseudomonas Colonization

Glyphosate -41%

Vitazyme +8%

Vitazyme + Glyphosate... +33%

Changes in IAA-Producing Bacteria Colonization

Glyphosate -29%

Vitazyme +20%




Vitazyme + Glyphosate... +42%

Glyphosate increased Fusarium infection, but Vitazyme combined with glyphosate reduced the infection below both Vitazyme and the control.

Glyphosate reduced Fluorescent Pseudomonas colonization by 41% below the control, while Vitazyme + Glyphosate increased levels well above the control and Vitazyme levels.

Glyphosate reduced IAA-producing bacteria below the control, while Vitazyme increased the levels, and the combined products increased levels by 42% above the control.

Vitazyme Effects on Fusarium and Bacterial Organism Colonization



Changes with Vitazyme




Fusarium -16%

Fluorescent Pseudomonas ... +51%

IAA-Producing Bacteria 53%

All Vitazyme treated plots were compared with all untreated plots, which revealed a highly significant reduction in Fusarium with Vitazyme, while beneficial bacteria increased by significant 51% and 53% levels.

Glyphosate Effects on Fusarium and Bacterial Organism Colonization



Changes with Glyphosate

Fusarium +6%
Fluorescent Pseudomonas -7%
IAA-Producing Bacteria -3%

All glyphosate treated plots were compared with all untreated plots, which revealed a trend toward more *Fusarium* lesions and fewer beneficial bacteria, though none of these differences are significant.

Fatty Acids, Oxidizers, Reducers, Glyphosate, and Enzyme Levels in Rhizosphere Soil

Treatment	TPLFA ¹ nm/g soil	Mn Oxidizers ² cfu/g soil x10 ⁴	Mn Reducers ³ cfu/g soil x10 ⁴	Red/Ox Ratio ⁴ cfu/g soil x10 ⁴	Glyphosate ⁵ ug/kg soil	B-Glucosamine Activity ⁶ ug PNP/g soil/hr
1. Control	133	6.25	8.47	1.38	274	173
2. Glyphosate	105 (-21%)	11.9 (+90%)	6.08 (-28%)	0.604 (-56)	940 (+243%)	147 (-15%)
3. Vitazyme	121 (-9%)	4.25 (-32%)	8.85 (+4%)	2.09 (+51%)	501 (+83%)	262 (+51%)
4. Vitazyme + Glyphosate	122 (-8%)	5.66 (-9%)	8.97 (+6%)	1.64 (+19%)	1,000 (+265%)	216 (+25%)
P>F	0.077	0.154	0.334	0.391	0.383	0.683

¹TPLFA = total phospholipid fatty acids; ²manganese oxidizing bacteria; ³manganese reducing bacteria; ⁴ratio of manganese reducing to oxidizing bacteria; ⁵soil glyphosate levels; ⁶soil level of this enzyme.

Vitazyme Effects on Fatty Acids, Oxidizers, Reducers, Glyphosate, and Enzymes

Treatment	TPLFA nm/g soil	Mn Oxidizers cfu/g soil x10 ⁴	Mn Reducers cfu/g soil x10 ⁴	Red/Ox Ratio cfu/g soil x10 ⁴	Glyphosate ug/kg soil	B-Glucosamine Activity ug PNP/g soil/hr
1. Control	119	9.07*	7.27	0.991	607	160
2. Vitazyme	122	5.04	8.91	1.86**	750	239**
P>F	0.718	0.019	0.218	0.001	0.148	0.009

*Significantly greater than the control at P=0.05; **Significantly greater than the control at P=0.01.

All Vitazyme treated plots were compared with all untreated plots. Note that manganese oxidizers were significantly reduced with Vitazyme, while manganese reducers increased. The reduced form of Mn is the form utilized by plants. This change in

oxidation state for Mn resulted in a highly significant increased in the Red/Ox Ratio with Vitazyme. The soil enzyme level was also significantly increased with Vitazyme.

Glyphosate Effects on Fatty Acids, Oxidizers, Reducers, Glyphosate, and Enzymes

Treatment	TPLFA nm/g soil	Mn Oxidizers cfu/g soil x10 ⁴	Mn Reducers cfu/g soil x10 ⁴	Red/Ox Ratio cfu/g soil x10 ⁴	Glyphosate ug/kg soil	B-Glucosamine Activity ug PNP/g soil/hr
1. Control	127	5.33	8.66	1.73	387	218
2. Glyphosate	113 (-11%)	8.77* (+65%)	7.52 (-13%)	1.12*** (-35)	970*** (+151)	181 (-17%)
P>F	0.083	0.038	0.380	0.008	0.001	0.158

*Significantly greater than the control at P=0.001.. **Significantly different than the other treatment at P=0.01.

All glyphosate treated plots were compared with all untreated plots. Phospholipids and soil enzymes were reduced by glyphosate, as were Mn reducers while Mn oxidizers were significantly increased (+65%) above the control. The Red/Ox Ratio was reduced by 35%, meaning few Mn⁺⁺ ions were available with glyphosate, and glyphosate residues were increased by an amazing 151% with the glyphosate treatment.

Conclusions: This University of Missouri small-plot corn trial, using a Vitazyme in-furrow and foliar application, with and without glyphosate for the foliar application, and with glyphosate alone, revealed that Vitazyme remediated the negative effects of the herbicide on soil organisms and other parameters. Yield was increased by 5 bu/acre — as for glyphosate — and Fusarium root infection was significantly decreased by 16%; at the same time, Fluorescent Pseudomonas and IAA-producing bacteria

were significantly increased. Manganese-reducing bacteria were increased by Vitazyme, and Mn-oxidizing bacteria were significantly reduced with Vitazyme, while soil B-glucosamine activity was increased as well. Glyphosate, on the other hand, produced opposite effects from Vitazyme by increasing Mn-oxidizers and Fusarium infection, while reducing beneficial microorganisms like Fluorescent Pseudomonas, IAA-producing bacteria, and Mn-reducers, while reducing phospholipid fatty acids and soil enzyme activity. Most notably, when Vitazyme was added to the glyphosate before spraying, the negative effect of the glyphosate on Fusarium and beneficial bacteria were reversed. These data indicate that, besides being a yield booster, Vitazyme, when applied with glyphosate herbicide, can remediate the damage this herbicide can do to the plant and soil, and thus is a natural remedy for farmers to add along with glyphosate in the sprayer tank.



Corn with Vitazyme application

Researcher: V. V. Plotnikov

Research institution: Agro Expert International, Vinnytsya, Ukraine

Location: L. L. C. Zlahoda, Vyshnivtsi Village, Onuphriy District, Kirovohrad Region, Ukraine

Variety: DKS 3511, FAO 330

Planting date: April 29, 2016

Seeding rate: 70,000 seeds/ha

Previous crop: winter wheat

Soil type: dark gray podzolic (2.6% organic matter)

Cultivation: disking to 6-8 cm, plowing to 20-22 cm, harrowing, and cultivation to 4-5 cm

Rainfall: 500-550 mm

Experimental design: A corn field was divided into a Vitazyme treated and untreated area, with the objective of determining the effect of this product on the yield of the grain.

① Control ② Vitazyme

Fertilization: 115 kg/ha of N pre-plant; at planting instead of 12-52-0 kg/ha of N-P₂O₅-K₂O in-row

Vitazyme application: 1.0 liter/ha on the leaves and soil at the 7 to 8-leaf stage on June 2, 2016

Yield results:

Treatment	Grain yield tons/ha	Yield change tons/ha
Control	4.8	—
Vitazyme	6.4	1.6 (+33%)


Increase in grain yield with Vitazyme: 33%

Income results: Vitazyme increased net profit by 263 USD/ha.

Conclusions: A corn trial in central Ukraine using Vitazyme at 1.0 liter/ha on the leaves and soil produced a 33%

yield increase. Profits were substantially increased proving the viability of this product for corn production in central Ukraine. A yield increase of this magnitude resulting from the 127 kg/ha of N used in this trial reveals the improvement in nitrogen efficiency this program affords.

Grain yield





Corn with Vitazyme application

Researcher: V. V. Plotnikov
Research institution: Agro Expert International, Vinnytsya, Ukraine
Location: Family Enterprise Kolyvailo, Miziakivs'ki Hutory Village, Vinnytsya Region, Ukraine.

Variety: DKS 4408 FAO 340
Planting date: April 25, 2016
Seeding rate: 75,000 seeds/ha
Previous crop: winter wheat
Soil type: Chernozem (3.0% organic matter)

Cultivation: disking to 6-8 cm, plowing to 20-22 cm, harrowing, and cultivation to 4-5 cm

Rainfall: 500-550 mm

Experimental design: A corn field was divided into a Vitazyme treated and untreated area, with the objective of determining the effect of this product on the yield of the grain.

Fertilization: 92 kg/ha of N pre-plant; at planting, 10-26-26 kg/ha of N-P₂O₅-K₂O in-row; 46 kg/ha of N side-dressed


Vitazyme application: 1.0 liter/ha on the leaves and soil at the 7 to 8-leaf stage on May 31, 2016

① Control ② Vitazyme

Yield results:

Treatment	Grain yield tons/ha	Yield change tons/ha
Control	11.1	—
Vitazyme	12.8	1.7 (+15%)

Increase in grain yield with Vitazyme: 15%



Income results: Vitazyme increased net profit by 281 USD/ha.

Conclusions: This Vitazyme foliar and soil treatment trial in Ukraine showed that only 1 liter/ha produced a 15% yield increase. Profits were also substantially increased, showing the viability of this product for corn production in central Ukraine on Chernozem soils.



Corn with Vitazyme application—A Nitrogen Efficiency Study



Notice the much improved root development when Vitazyme is applied to the seeds and leaves.



The research plots at South Dakota State University produced excellent yield and nitrogen efficiency data for 2016.

Researcher: Dr. David Clay and Graig Reicks

Research institution: Department of Plant Sciences, South Dakota State University, Brookings, South Dakota

Location: South Dakota State University, Field Research Facility, Aurora, South Dakota

Variety: DKC 49-72 (99-day hybrid)

Planting date: May 14, 2016

Planting rate: 32,000 seeds/acre

Planting depth: 2 inches

Row spacing: 30 inches

Soil type: Strayhoss-Maddox Complex, 2% slope, loam texture

Experimental design: A replicated randomized complete block design, with plots 6 rows wide (15 feet) and 20

feet long, was established to determine the effectiveness of Vitazyme as both a seed and a soil treatment for corn. Yield measurements were made, and also ¹⁵N measurements were made using a mass spectrometer to determine the efficiency of nitrogen use within the plant. Three fertilizer nitrogen rates were used, and four replications.

Fertilization: Nitrogen fertilizer was applied by hand as urea (46%) to appropriate plots on June 13, 2016, when the corn was at the V3 stage.

Vitazyme application: Seed treatment (at planting on May 14): 112 ml of Vitazyme per 10 lb of seed, or 7.4 oz/acre sprayed on the seeds and mixed on a tarp; **Foliar treatment** (on June 29 at V 8-9) at 13 oz/acre, using a 15 gal/acre spray solution, with a non-ionic surfactant (0.25% v/v) and a backpack sprayer.

¹⁵N determinations: A mass spectrometer was used at the Department of Plant Science, South Dakota State University.

Yield and nitrogen results: At the time of this reporting, only the seed treatment results are given. The full report will be available at a later date.


Treatment	Nitrogen fertilizer	Corn yield ¹	Corn grain use efficiency ²
	bu/acre	bu/acre	%
1. None	125	220 a	43.3 b
	75	197 b	40.3 b
	0	181 bc	—
2. Vitazyme on seeds	125	218 a	42.6 b
	75	218 a	58.0 b
	0	162 c	

¹Means followed by the same letter are not significantly at P=0.05.

²Means followed by the same letter are not significantly different at P=0.10. The calculations used to determine this value are based upon 14N and 15N ratios in the grain, and will not be presented in this report.

Nitrogen rate	Vitazyme application			
	None	Seeds	Leaves	Seeds + Leaves
lb/acre	(treatment number)			
0	1	4	7	10
75	2	5	8	11
125	3	6	9	12


Vitazyme Seed Treatment Effect on Corn Yield



Increase in corn yield with Vitazyme seed treatment: 21 bu/acre, or 11%

At the same nitrogen level, with or without a Vitazyme seed treatment, the yield increased by 11%, which was the same yield as for the high nitrogen rate (125 lb/acre) with the seed treatment. Thus, the nitrogen utilization was improved greatly with Vitazyme, as evidenced by the Corn Grain Use Efficiency value.

Vitazyme Seed Treatment Effect on Corn Grain Use Efficiency at 75 lb/acre nitrogen



Ear and root development are improved with a seed and a foliar Vitazyme treatment, as seen here.

Improvement in nitrogen use efficiency of the grain with Vitazyme on seeds: 17.7 %-points

This improvement in the use of nitrogen by the seeds was dramatically shown at the reduced N rate of 75 lb/acre. It did not occur at the 125 lb/acre N rate, showing the potential of Vitazyme's active agents to greatly improve N transfer into the grain when fertilizer N is somewhat limiting.

Conclusions: A small plot, replicated corn trial near Brookings, South Dakota, using 12 treatments with a Vitazyme seed treatment, a foliar treatment, or both, at three nitrogen levels (0, 75, and 125 lb of N/acre), revealed a pronounced yield response at the reduced nitrogen level (75 lb/acre) for the seed treatment. The corn yield jumped by 11%, from 197 to 218 bu/acre, a highly significant increase

that equalled the seed treatment yield at the high nitrogen level (125 lb/acre). This high yield was about the same as the yield obtained at 125 lb/acre of nitrogen without Vitazyme on the seeds. Calculations of nitrogen use efficiency using ¹⁵N computations revealed that, at the reduced 75 lb/acre nitrogen rate, the efficiency of nitrogen movement into the grain was greatly enhanced with the

Vitazyme seed treatment, increasing from 40.3% to 58.0%. This increase occurred at the reduced nitrogen rate only, revealing that when more fertility stress existed the brassinosteroids and other active agents in Vitazyme caused more nutrients to be translocated into the grain out of the total fertilizer added, compared to the untreated corn at the same 75 lb/acre nitrogen rate.

Corn South Dakota State University—Nitrogen and Water Use Efficiency



Corn receiving 125 lb/acre of N in this nitrogen efficiency trial at South Dakota State University produced a much greater root mass when treated with Vitazyme.



Typical ears for the 125 lb/acre nitrogen treatment reveal greater ear size with Vitazyme applied twice, leading to a 9 bu/acre (6%) yield increase and better nitrogen utilization.

Researchers: David Clay, Ph.D., and Craig Reicks, Research Assistant II

Research institution: South Dakota State University, Department of Plant Science, Brookings, South Dakota

Location: Aurora, South Dakota

Variety: NK N41Y-3000 GT (98-day hybrid)

Plant population: 32,000 seeds/acre

Tillage: conventional

Soil type: Brandt silty clay loam

Planting date: May 16, 2014

Experimental design: A site for small plot studies was selected at the South Dakota State University research facility near Aurora. Plots of 10 x 20 feet (5 x 10 feet harvested) were laid out with four treatments in a randomized complete block design to determine the effects of Vitazyme, applied to the seed, leaves, and both, on the yield, nitrogen-use efficiency, and water-use efficiency of corn. Three nitrogen rates were used.

Nitrogen rate lb/acre	Vitazyme application			
	None	Seeds	Foliage	Seeds + Foliage
	(treatment number)			
0	x(1)	x(4)	x(7)	x(10)
75	x(2)	x(5)	x(8)	x(11)
125	x(3)	x(6)	x(9)	x(12)

Fertilization: Urea was broadcast on June 4 over the appropriate plots at 75 or 125 lb/acre of nitrogen around planting time.

Vitazyme application: (1) seed treatment of 5% Vitazyme, with 3.2 oz misted and mixed by hand with 10 lb of seed; (2) 13 oz/acre sprayed on the leaves and soil at the 6-leaf stage (V6) on July 9, at 15 gal/acre of solution. A non-ionic surfactant was added at 0.5% of the total solution.

Growing season weather: wetter than normal (12.33 inches for June through August, versus the average of 10.73 inches), and cooler than normal (1,585 growing degree days versus the average of 1,668)

Harvest date: The middle two rows of each plot were hand-picked on October 31, ears were dried at 140° F, and then shelled and weighed.


Yield results: NOTE: There were planting difficulties for the seeds treated with Vitazyme—populations were low—so these plots were replanted once the error was detected. However, the plants failed to respond normally, so these data have been eliminated from some analyses.

Nitrogen Fertilizer Effects

Treatment	Yield ¹ bu/acre	Yield change bu/acre
No nitrogen	106 c	—
75 lb/acre nitrogen	146 b	40 (+38%)
125 lb/acre nitrogen	160 a	54 (+51%)

¹ Yields followed by different letters are significantly different at P=0.01.

Nitrogen fertilizer produced a straight-line yield response over the three N levels, and over all Vitazyme treatments.



Vitazyme Effects at Three Nitrogen Levels


In this analysis, plots that received and did not receive the Vitazyme seed treatment are included.

Vitazyme applied to both the seeds and foliage overcame the drop in corn yield with the seed treatment above. The reason for this is not understood.

Corn South Dakota State University—Nitrogen and Water Use Efficiency cont.

Nitrogen rate lb/acre	Vitazyme	Grain yield bu/acre	Yield change bu/acre
0	0 Foliage	97 116	— +19 ^a (+20%)
75	0 Foliage	144 148	— +4 ^b (+3%)
125	0 Foliage	155 164	— +9 ^c (+6%)

^aSignificant at P=0.09; ^bSignificant at P=0.40; ^cSignificant at P=0.03.





Yield, nitrogen efficiency, and water efficiency effects:

The determination of nitrogen (N) and water efficiency is done through the calculation of isotopes of nitrogen and carbon, using a mass spectrometer. The calculation of these efficiencies, and related parameters, is complex and will not be discussed here, but can be reviewed in a paper by K. Kim, D. Clay, C. Carlson, S. Clay, and T. Trooien entitled "Do synergistic relationships between nitrogen and water influence the ability of corn to use nitrogen derived from fertilizer and soil?" (Agronomy Journal 100 [3], 2008, pages 551-557).

The following analyses were made using only the foliar Vitazyme applications, to avoid the problems associated with the seed treatment, as discussed earlier. Probabilities of significant differences are indicated beneath each pair of values.

N-rate	Treatment	N-efficiency bu/lb of N	N-balance ¹	¹⁵ N%	¹³ C%	¹⁵ N%	¹³ C%	Soil inorganic N balance lb/acre
				—% in grain—		—% in stover—		
0	Vitazyme	—	- 65.0	0.36	- 11.2	- 0.83	- 11.9	0.23
	Control	—	- 85.0	1.05	- 11.3	- 2.81	- 11.8	- 11.2
			p=0.48	p=0.41	p=0.72	p=0.30	p=0.34	p=0.58
75	Vitazyme	0.538	- 42.0	0.116	- 11.19	- 2.12	- 11.93	- 5.6
	Control	0.563	- 64.2	0.632	- 11.26	- 0.53	- 11.97	- 14.3
		p=0.61	p=0.11	p=0.16	p=0.08	p=0.65	p=0.79	p=0.49
125	Vitazyme	0.466	19.6	0.765	- 11.33	- 0.66	- 11.94	29.5
	Control	0.400	- 20.5	1.251	- 11.43	5.76	- 12.11	- 16.2
		p=0.01	p=0.01	p=0.44	p=0.23	p=0.02	p=0.43	p=0.01

¹A higher value means more efficient uptake of N.



Increase in N-efficiency with Vitazyme: 17%


At all three nitrogen rates, foliar Vitazyme treatment increased N-use efficiency, especially at 125 lb/acre N. At the typical farm-applied rate of 125 lb/acre of N, Vitazyme treated corn produced 0.066 more bushels/acre per pound of nitrogen than the control.

Corn South Dakota State University—Nitrogen and Water Use Efficiency cont.


N-rate	Treatment	Grain yield	Yield loss, nitrogen stress	Yield loss, water stress
lb/acre		bu/acre	bu/acre	bu/acre
0	Vitazyme	94.4	20.1	60.9
	Control	105.8	12.4	57.2
		p=0.49	p=0.32	p=0.75
75	Vitazyme	146.2	5.66	23.7
	Control	152.0	2.60	21.1
		p=0.16	p=0.58	p=0.75
125	Vitazyme	170.1	0.634	5.0
	Control	161.0	0.470	14.2
		p=0.01	p=0.53	p=0.02

Conclusion: A corn trial in eastern South Dakota designed to evaluate the effects of Vitazyme seed and foliar treatments on grain yield, and nitrogen (N) and water use efficiency at three N levels, was impaired by a poor plant population for the seed treated plots. Thus, in most cases only the foliar treatment treated at V6 was evaluated. In spite of this limitation, Vitazyme improved grain yield significantly at the 125 lb/acre N rate (+6%), increased N-efficiency by 17%, and reduced the yield loss due to water stress by 9.2 bu/acre; the control yield loss at 125 lb/acre N was 14.2 bu/acre. The 0 and 75 lb/acre N rates did not show significant yield or N and water use responses, for unknown reasons. Moreover, N-balance was improved at all fertility levels, especially at the 125 lb/acre N application rate. Since this highest of the three rates is similar to a typical farmer N application rate, the value of Vitazyme use for improved nitrogen and water use efficiency is displayed in this study, the result being a 9.1 bu/acre yield increase. A seed treatment along with the foliar application would likely have triggered significant responses for all parameters at all three N application rates.

Grain Yield with 125 lb/acre N



Yield Loss from Water Stress at 125 lb/acre N



Corn with Vitazyme application

Researchers: Ernesto Infante, Lucero Fernandez, and Edgar Ortiz

Research organizations: La Mazorca and Quimica Lucava

Location: Daniel Hernandez Farm, La Barca, Jalisco, Mexico

Variety: Asgrow Antelope

Planting date: April 30, 2015

Row Spacing: 0.75 meter

Experimental design: A corn field of 4.75 ha was selected to treat 1.0 ha with Vitazyme, to determine effects of the product on crop growth, insect infestation, and yield. Two Vitazyme applications were made.

1 Control 2 Vitazyme


Fertilization: unknown

Vitazyme application: (1) seed treatment on a tarp before planting on April 30 using 250 ml of Vitazyme with a backpack sprayer, and allowing the seeds to dry before planting; (2) 1 liter/ha sprayed on the leaves and soil 30 days later on May 30

Harvest date: October 20, 2015

Growth results: The treated corn grew much faster than the control, being 10 to 15 cm taller in the early stages with darker green leaves and considerably less incidence of fall army worm. A growth difference was visible even up to harvest, when the Vitazyme treated portion of the field was taller, with more leaf area, and larger ears.

Yield results: On October 20, 16 rows of the treated and the control areas were harvested for 205.6m, giving a harvested area of 2,467.2m²



Treatment	Plot yield kg/2,467m ²	Grain moisture %	Moisture-corrected yield ¹ kg/ha	Yield change kg/ha
Control	3,385	13.96	13,726.4	—
Vitazyme	3,670	13.70	14,927.1	1,200.7 (+9%)

¹Correction factor: (100-% H₂O)/86; Dry yield = (Correction factor) (Harvested yield/ha).

Increase in corn yield with Vitazyme: 9%

Income results: Using a corn price of \$0.225 (U.S.)/kg, the income is as follows:

Treatment	Dry yield tons/ha	Gross income USD/ha	Income change USD/ha	Net income change ¹ USD/ha	Cost : benefit ²
Control	13.7264	3,088.44	—	—	—
Vitazyme	14.9271	3,358.60	270.16	226.89	5.2

¹Vitazyme costs for overall 1.25 L/ha, including labor = 43.34 USD/ha. ²Cost : Benefit = (Net income increase)/(Vitazyme cost).

Conclusions: This corn study in Mexico revealed that a 0.25 liter/ha Vitazyme seed treatment at planting, followed by a 1.0 liter/ha foliar and soil application 30 days later, greatly improved corn growth and yield. Faster growth of the plant, more leaf chlorophyll, and less army worm incidence led to a yield increase of 9%, and an excellent cost : benefit of 5.2, showing the excellent efficacy of this program for corn growers in Mexico.

Corn with Vitazyme application

Researchers: Bartolo González and Lucero Fernandez

Farmer: Irrigation District 011

Research organizations: CVYTTS and Quimica Lucava

Location: Irapuato, Guanajuato, Mexico

Variety: CERES XR47

Planting date: May 12, 2014

Experimental design: A 1 hectare area of a corn field was treated twice with Vitazyme to determine its effects on growth, yield, and profitability compared to an adjoining untreated control area.

Fertilization: unknown

Vitazyme application: (1) a seed treatment at 250 ml/ha on May 12, 2014; (2) foliar and soil spray at 1 liter/ha on June 20, 2014

Growth observation: The Vitazyme treated plants had *more extensive roots* and *greater plant vigor*.


Harvest date: December 16, 2014

Yield results:

Treatment	Yield	Yield change
	kg/ha	kg/ha
Control	9,670	—
Vitazyme	10,437	767 (+8%)

Increase in corn yield with Vitazyme: 8%

Income results: The corn price was USD 0.25/kg. Vitazyme overall cost, including labor = USD 46.43/ha.



Added income with Vitazyme: 191.75 USD/ha

Added profit with Vitazyme: 145.32 USD/ha

Cost : Benefit with Vitazyme: 3.13

Conclusions: This Mexican corn trial showed an excellent yield increase (8%) and income improvement (145.32 USD/ha) with two Vitazyme applications, proving it's great efficacy for use in corn production in Mexico.

Corn with Vitazyme application

Researcher: V. V. Plotnikov

Research organizations: Agricultural LLC, "Palmira Vidhodyvlya", Ukraine

Location: Zolotonosha District, Cherkasy Region, Voznesens'ke Village, Ukraine

Variety: DKS 4590 hybrid

Seeding rate: 75,000/ha

Planting date: April 26, 2015

Previous crop: soybeans

Cultivation methods: disking to 8 cm, plowing to 24 cm, cultivation to 6 cm

Experimental design: A trial with Vitazyme on corn involved dividing a field into a Vitazyme treated and an untreated area. The purpose of the trial was to evaluate the product's effects on corn yield and profitability.

Fertilization: 10 kg/ha N, 26 kg/ha P₂O₅, and 26 kg/ha K₂O in-furrow at planting; 115 kg/ha N broadcast pre-plant

Vitazyme application: 1 liter/ha on the leaves and soil on May 31, 2015, at the 7 to 8-leaf stage


Yield results: Harvest date is unknown

Treatment	Yield tons/ha	Yield change tons/ha
Control	10.2	—
Vitazyme	11.8	1.6 (+16%)

Increase in grain yield with Vitazyme: 16%

Income results: A 1 liter/ha application gave a profit increase of 4,710 UAH/ha.

Conclusions: In this Ukraine corn trial, a single soil and foliar Vitazyme application of 1 liter/ha, at the 7 to 8-leaf stage, resulted in a substantial yield increase of 16%, and a profit improvement of 4,710 UAH/ha. This result indicates the great efficacy of this program for use by corn growers in Ukraine.



Corn with Vitazyme application

Researcher: V. V. Plotnikov

Research organization: Small Private Enterprise, "Firm Harant", Ukraine

Location: Teplyts'kyi District, Vinnytsya Region, Teplyk Town, Ukraine

Variety: P9175 hybrid

Seeding rate: 70,000/ha

Planting date: April 24, 2015

Previous crop: winter wheat

Soil type: podzolized chernozem

Cultivation methods: disking to 8 cm, plowing to 24 cm, cultivation to 6 cm

Experimental design: A corn field was divided into a Vitazyme treated and an untreated area to determine the effectiveness of Vitazyme for improving grain yield and profitability.

1 Control 2 Vitazyme

Fertilization: 18 kg/ha N, 18 kg/ha P₂O₅, and 18 kg/ha K₂O in-furrow at planting;

90 kg/ha N pre-plant broadcast

Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 7 to 8-leaf stage, on May 28, 2015.


Yield results: Harvest date is unknown

Treatment	Yield	Yield change
	tons/ha	tons/ha
Control	7.14	—
Vitazyme	8.35	1.21 (+17%)

Increase in corn yield with Vitazyme: 17%

Income results: Profit increased by 3,442 U UAH/ha with Vitazyme.

Conclusions: A corn trial in Ukraine in 2015, with 1 liter/ha sprayed on the leaves and soil at the 7 to 8-leaf stage, resulted in the yield increasing by a substantial 17%, giving 3,442 UAH/ha more profit. This great increase reveals the considerable efficacy of this program for corn in Ukraine.



Corn *A Synergism Study with Seaweed*

Researcher: Bertel Schou, Ph.D.

Research organization: ACRES

Research, Cedar Falls, Iowa

Location: Cedar Falls, Iowa

Variety: Pioneer P0636AM

Seeding rate: 38,000 seeds/acre

Planting depth: 1.5 inches

Row width: 30 inches

Soil type: Floyd loam (pH=6.3, organic matter=4.0%, cation exchange capacity=15.5 meg/100g, fertility level=excellent, drainage=excellent)

Planting date: May 2, 2015

Experimental design: A small-plot corn trial, using a randomized and complete block design and four replicates, was laid out in plots that were 15 (six rows) x 30 feet, with the objective of determining the effectiveness of Vitazyme and seaweed to influence corn yield, alone and together.



Corn treated twice with Vitazyme in this Iowa trial produced a much more massive root system, plus a 5% yield increase, than the untreated control. Seaweed did not produce a synergism with Vitazyme.

- 1 Control
- 2 Vitazyme
- 3 Seaweed
- 4 Vitazyme + Seaweed

Fertilization: All plots received a fall (2014) broadcast application of 18-16-60 16/acre of N-P₂O₅-K₂O, and 100 lb/acre of N in 2015.

Vitazyme application: For Treatments 2 and 4, (1) 13 oz/acre in-furrow at planting on May 2; (2) 13 oz/acre on the leaves and soil at V6 on June 24.

Seaweed application: For Treatments 3 and 4, (1) 2 qts./acre in-furrow at planting on May 2; (2) 2 qts./acre on the leaves and soil at V6 on July 2; (3) 2

qts./acre on the leaves and soil at VT on July 29. The seaweed was from Ocean Organics bared in Waldoboro, Maine, and Ann Arbor, Michigan.

Growing season weather: Excellent, with rainfall evenly distributed throughout the spring and summer, and average temperatures were good for growth. Rainfall; April = 2.97 in.; May=5.62 in.; June=5.40 in.; July=4.65 in.; August=7.50 in.; September=3.33 in. Temperature (daily average maximum): April=66°, May=70°; June=80°, July=85°, August=79°, September=81°.

Harvest date: October 10, 2015

Corn moisture and population results:

Grain moisture varied from 16.08 to 16.75% among treatments, and was not significant, while plant population also varied little among treatments.


Yield results: The corn was harvested using a Massey-Ferguson plot combine, with an electronic scale and moisture meter.

Increase in Corn Yield

- Vitazyme alone 5%
- Seaweed alone 4%
- Vitazyme + Seaweed 2%

Treatment	Corn yield ¹ bu/acre	Yield change bu/acre
1. Control	215.78 b	—
2 Vitazyme	225.58 a	9.80 (+5%)
3. Seaweed	224.38 ab	8.60 (+4%)
4. Vita + Seaweed	220.70 ab	4.92 (+2%)
LSD (P=0.05) (P ₂ 0.05)	9.52	
CV	2.79	
Replicate probability	0.6926	
Treatment probability	0.2542	

¹ Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.



Corn *A Synergism Study with Seaweed cont.*

Conclusions: A corn study conducted in east-central Iowa, using small plots with four replications, revealed that Vitazyme, applied at planting in-furrow and sprayed at V6, significantly increased corn yield by 9.80 bu/acre (5%). Seaweed alone, applied in-furrow and twice foliar, increased the yield by 8.60 bu/acre (4%), but the two products combined did not provide additive effects, increasing the yield by 4.92 bu/acre (2%). Grain moisture and plant population were not significantly affected by either product. Results in 2014 with corn at ACRES Research did not show a synergism between the two products when they were applied together, and it was hoped that by separating foliar applications by several days a synergism might occur. However, it did not. Both Vitazyme and seaweed applied alone gave excellent responses to corn yield in 2015.

Corn *A Study with Wet-Sol Adjuvant*

Researcher: Bertel Schou, Ph.D.

Research institution: ACRES (Agricultural Research and Education Services), Cedar Falls, Iowa

Location: Cedar Falls, Iowa

Variety: Pioneer P0636AM (GMO-RR)

Planting date: April 30, 2015

Planting rate: 38,000 seeds/acre

Planting depth: 1.5 inches

Row spacing: 30 inches

Soil type: Floyd Loam (pH=6.1, organic matter=4.0%, cation exchange capacity=15.5 meg/100g, fertility level=excellent, drainage=excellent)

Experimental design: A small-plot corn study was arranged in a randomized complete block design and four replicates, with plots that were 15 (6 rows) x 30 feet; the two center rows from each plot were harvested. The purpose of the study was to evaluate the effect of Wet-Sol 233, a non-ionic adjuvant, at different rates in-furrow and foliar, with and without Vitazyme, on the growth and yield of corn.



The dramatic improvement in rooting with Vitazyme can be seen on the right; two applications improved yield by 13.1 bu/acre (6%) above the untreated control.

Treatment	Vitazyme		Wet-Sol 233	
	In-furrow	Foliar	In-furrow	Foliar
-----oz/acre-----				
1	0	0	0	0
2	13	13	0	0
3	0	0	16	16
4	0	0	32	16
5	0	0	48	16
6	13	13	16	16

Fertilization: All plots received 18-16-60 lb/acre of N-P₂O₅-K₂O in the fall of 2014, and 1.00 lb/acre of N in 2015.

Vitazyme application: For Treatments 2 and 6, (1) 13 oz/acre in-furrow at planting on April 30, 2015; (2) 13 oz/acre on the leaves and soil at V7 on June 23, 2015.

Wet-Sol 233 application: Wet-Sol 233 is a non-ionic surfactant produced by Scheaffer Oil, St. Louis, Missouri. For Treatments 3 to 6, (1) 16 to 48 oz/acre in-furrow at planting on April 30, 2015; (2) 16 oz/acre on the leaves and soil at V7 on June 23, 2015. For Treatment 6, Vitazyme was mixed with Wet-Sol.

Growing season weather: Excellent, with rainfall evenly distributed throughout the spring and summer, and the average temperatures were good for growth. Rainfall: April=2.97in.; May=5.62 in.; June=5.40 in.; July=4.65 in.; August=7.50 in.; September=3.33 in, Temperature (daily average maximum): April=66; May=70; June=80; July=85; August=79; September=81.

Harvest date: October 8, 2015

Plant population results (final): There were no significant differences among treatments from final population.

Grain moisture results: Harvest was completed using a Massey-Ferguson plot combine having an electronic scale and moisture tester.


Treatment	Grain moisture ¹	Moisture change
	%	%
1. Control	16.663 a	—
2. Vitazyme	16.655 a	(-)0.008
3. Wet-Sol 16 oz/A	15.965 a	(-) 0.698
4. Wet-Sol 32 oz/A	15.743 a	(-) 0.920
5. Wet-Sol 48 oz/A	16.363 a	(-) 0.300
6. Vita + Wet-Sol	16.063 a	(-) 0.600
LSD (P=0.05)	0.941	
CV	3.84	
Replicate probability	0.0334	
Treatment probability	0.2548	

¹ Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.

While none of the treatment means are significant, there was a definite trend for all of the Wet-Sol treatments to produce drier grain at harvest, up to 0.92% point at 32 oz/acre; Vitazyme alone did not produce such a decrease.

Corn *A Study with Wet-Sol Adjuvant cont.*

Corn Grain Moisture



Conclusion: A corn study in east-central Iowa in 2015, a very good cropping year with near-record yields in the area, showed that both Vitazyme and Wet-Sol 233 produced significantly greater yields ($P=0.05$) than the control. This increase occurred at the 16, 32, and 48 oz/acre rates, with Vitazyme alone producing the highest overall yield of 223.0 bu/acre, an increase over the control of 6%. Vitazyme combined with Wet-Sol 233 did not produce an additive effect on yield, with a 4% yield increase. Wet-Sol 233 tended to produce drier corn at harvest, though not significantly so. These results show the ability of both Vitazyme and Wet-Sol to increase corn grain yields during an especially good cropping year, and their viability as major production tools for agriculture in the Corn Belt of the United States.

Decrease in Grain Moisture, %-points

Wet-Sol, 16 oz/acre.....	0.698
Wet-Sol, 32 oz/acre.....	0.920
Vitazyme + Wet-Sol, 16 oz/acre	0.600


Yield results:

Treatment	Corn Yield ¹ bu/acre	Yield change bu/acre
1. Control	209.9 b	—
2. Vitazyme	223.0 a	13.1 (+6%)
3. Wet-Sol 16 oz/A	221.9 a	12.0(+6%)
4. Wet-Sol 32 oz/A	221.1 a	11.2 (+5%)
5. Wet-Sol 48 oz/A	221.6 a	11.7 (+6%)
6. Vita + Wet-Sol	217.3 ab	7.4 (+4%)
LSD (P=0.05)	8.2	
CV	2.49	
Replicate probability	0.0001	
Treatment probability	0.0355	

¹ Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.

Vitazyme alone produced the greatest yield increase, at 13.1 bu/acre (6%), a significant increase above the control but statistically equal to all three Wet-Sol treatments. Vitazyme and Wet-Sol together did not produce a positive interaction, though the yield was not significantly less than the other treatments; neither was it greater than the control.

Corn Grain Moisture



Increase in Corn Yield, %

Vitazyme only	6%
Wet-Sol, 16 oz/acre.....	6%
Wet-Sol, 32 oz/acre.....	5%
Wet-Sol, 48 oz/acre.....	6%
Vitazyme + Wet-Sol, 16 oz/acre	4%

Corn *A Summary of Yield and Income Results in Mexico: 2012–2015*



Untreated corn in Atotonilco, Jalisco, Mexico, shows good growth, but nothing like that for Vitazyme (see the adjoining photo).



Two Vitazyme treatments greatly enhanced the growth of this corn, which is typical of the many trials conducted with this product in Mexico since 2012.

The following results for corn trials in Mexico were compiled by Juan Carlos Diaz of Ag Biotech.

Vitazyme application: Seed treatment at 0.25 liter/ha, and a foliar treatment at 1 liter/ha 30 days after planting

Conclusion: These eight Mexican corn trials revealed the excellent efficacy of Vitazyme use, with a seed treatment and single foliar application.

Test site	Year	Corn yield			Income ¹ USD/ha	Net Profit USD/ha	Cost : Benefit
		Control tons/ha	Vitazyme tons/ha	Change tons/ha			
El Monte, Villa Corona, Jalisco	2012	8.6	10.5	1.9 (+22%)	426.06	382.87	8.9
La Mesita, Cocula, Jalisco	2012	12.8	15.3	2.5 (+20%)	560.60	517.41	12.0
El Llano, San Martin Hidalgo, Jalisco	2012	7.30	8.21	0.91 (+13%)	204.06	160.87	3.7
Camino a la Coronilla, Arneco, Jalisco	2012	11.1	13.0	1.9 (+17%)	426.06	382.87	8.9
Odilón Ramos, San Juan Acozac, Puebla	2012	4.75	10.00	5.25 (+111%)	1,177.26	1,134.07	26.3
Las Margaritas, Ayotlan, Jalisco	2014	13.00	13.81	0.81 (+6%)	182.20	139.01	3.2
Distrito Riego 11 Irapuato, Guanajuato	2014	9.670	10.437	0.767 (+8%)	171.99	128.80	3.0
El Cabezón, Ameco, Jalisco	2014	11.51	12.84	1.33 (+12%)	297.57	254.38	5.9
Daniel Hernandez, La Barca, Jalisco	2015	13.73	14.93	1.20 (+9%)	269.24	226.05	5.2
Mean		10.27	12.11	1.84 (+18%)	412.78	369.59	8.6

¹ Corn price = 224.24 USD/ton. Vitazyme cost = 27.27 USD/liter; at 1.25 liters/ha, the total cost was 34.10 USD/ha. Spraying cost (labor) for two sprayings = 9.09 USD/ha. Overall Vitazyme cost = 43.19 USD/ha.

Yield increase with Vitazyme: +18%

Net profit increase with Vitazyme: 369.59 USD/ha

Cost: Benefit increase with Vitazyme: 8.6

It is clear that this program provides excellent yield and profit increases for corn farmers in Mexico.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2014 Crop Results

Vitazyme on Corn Five Trials in Mexico

Researcher: unknown

Research organization: Quimica Lucava

Experimental design: All trials utilized a split field, with Vitazyme applied to a portion to determine effects of the product on yield.

1. Control


2. Vitazyme (2X)

Fertilization: unknown

Vitazyme application: (1) 0.25 liter/ha on the seeds at planting; (2) 1 liter/ha sprayed on the plants and soil 30 days later.


El Monte, Jalisco, Mexico

Treatment	Corn yield	Yield change
	----- tons/ha -----	
Control	8.6	—
Vitazyme	10.5	1.9 (+22%)



Increase in corn yield with Vitazyme: 22%

La Mesita, Jalisco, Mexico




Treatment	Corn yield	Yield change
	----- tons/ha -----	
Control	12.8	—
Vitazyme	15.3	2.5 (+20%)

Increase in corn yield with Vitazyme: 20%


El Llano, Jalisco, Mexico

Treatment	Corn yield ----- tons/ha -----	Yield change
Control	7.30	—
Vitazyme	8.21	0.91 (+12%)

**Increase in corn yield
with Vitazyme: 12%**



Camino a la Coronilla, Jalisco, Mexico




Treatment	Corn yield ----- tons/ha -----	Yield change
Control	11.1	—
Vitazyme	13.0	1.9 (+17%)

**Increase in corn yield
with Vitazyme: 17%**

San Juan Acozac, Puebla, Mexico

Treatment	Corn yield ----- tons/ha -----	Yield change
Control	4.75	—
Vitazyme	10.00	5.25 (+111%)

**Increase in corn yield
with Vitazyme: 111%**



Economic Evaluation of All Trials

Income from the average yield increase (2.49 tons/ha)	7,476 pecos
Cost of Vitazyme (1.25 liters/ha)	500 pecos
Cost of application	150 pecos
Profit (income versus costs)	6,826 pecos
Cost:Benefit ratio	13.7:1

Conclusions: These five Mexican corn trials showed excellent yield responses to Vitazyme, when applied to the seeds and later to the leaves and soil. The average yield increase was 2.49 tons/ha, or 36% more than the control plants, giving an overall increase in profit of 6,826 pecos/ha (\$546.08/ha). With a cost:benefit rates of 13:7:1, this Vitazyme program is highly recommended for corn growers across Mexico.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2014 Crop Results

Vitazyme on Corn

University of Missouri - Bradford Research Center

Researcher: Manjula Nathan, Ph.D.
Row spacing: 30 inches

Location: University of Missouri Bradford Research Center,
Columbia, Missouri

Variety: Dekalb 62-97

Population: 32,000 seeds/acre

Planting date: May 16, 2013

Experimental design: A replicated corn study (six replications), in a randomized complete block design, was established using six-row plots for each treatment, each being 100 feet long (0.03443 acre/plot). The objective of the study was to evaluate the effect of Vitazyme, applied to the seeds and later on the leaves, on leaf nutrient and yield parameters.

1. Control

2. Vitazyme on the seeds and leaves


Fertilization: according to soil test recommendations using 160-46-62 lb/acre N-P₂O₅-K₂O before planting

Vitazyme application: (1) 13 oz/acre equivalent on the seeds, mixed thoroughly, just before planting (6.5 oz of Vitazyme applied to 16,000 seeds); (2) 13 oz/acre on the leaves and soil at the 8-leaf stage (knee height) on July 3.

Weather during the 2013 growing season: A wet spring delayed planting; otherwise, conditions for growth were favorable except for a short-duration drought in August and early September. Heavy rain after fertilizer application in May could have caused some nitrogen loss.


Tissue nutrient levels: Ear leaf samples were collected July 26 at silking. These samples were analyzed for N, P, K, Ca, and Mg at the University of Missouri.

Corn Tissue Nitrogen




P > F = 0.208

Corn Tissue Phosphorus



P > F = 0.604

Corn Tissue Potassium




P > F = 0.506

**Increase in N with
Vitazyme: 10%**

**Increase in P with
Vitazyme: 8%**


**Increase in K with
Vitazyme: 3%**

Corn Tissue Calcium



P > F = 0.034

Corn Tissue Magnesium



P > F = 0.079


Increase in Ca with Vitazyme: 39%

Increase in Mg with Vitazyme: 15%

Vitazyme increased all five tissue elements analyzed, with increases ranging from 3% (K) to 39% (Ca). The Ca and Mg increases were significant, especially Ca (39%), while N, P, and K increases were not significant.

Yield parameter results:


Grain Yield



P > F = 0.99. Two applications of Vitazyme greatly improved corn grain yield, by 11 bu/acre, above the control.


Increase in grain yield with Vitazyme: 7%

Ear Length




P > F = 0.345

Ear Circumference



P > F = 0.113

Rows On Ears




P > F = 0.317

Increase in ear length with Vitazyme: 5%

Increase in ear circumference with Vitazyme: 4%


Increase in rows on ears with Vitazyme: 0.9 row

Seeds Per Ear




P > F = 0.172

Weight Per Ear



P > F = 0.253

Weight Per Kernel



P > F = 0.474

Increase in seeds per ear with Vitazyme: 10%

Increase in ear weight with Vitazyme: 15%

Increase in weight per kernel with Vitazyme: 4%

Conclusions: A replicated corn trial at the University of Missouri in 2013 revealed that Vitazyme, applied on the seeds before planting, and again at 13 oz/acre to the leaves and soil at V8, improved tissue nutrient levels as well as yield and all harvest parameters, several significantly. A summary of these effects is as follows:

Increase with Vitazyme

Yield Parameters

Grain yield	10%*
Ear length	5%
Ear circumference	4%
Rows per ear	0.9 row/ear
Seeds per ear	10%
Weight per ear	15%
Weight per kernel	4%

Tissue Parameters

Nitrogen	10%
Phosphorus	8%
Potassium	3%
Calcium	39%*
Magnesium	15%*

*Significantly greater than the control, at P=0.10 or less.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2014 Crop Results

Vitazyme on Corn

Agricultural Custom Research and Education Services (ACRES)

Researcher: Bertel Schou, Ph.D.
Educational Services

Variety: Pioneer PO453 AM (Roundup Ready)

Planting date: May 8, 2014

Soil type: Aredale loam (36% sand, 42% silt, 22% clay, 4.6% organic matter, pH = 6.2, cation exchange capacity = 17.6 meq/100 g)

Soil test results: (Perry Agricultural Laboratory, Bowling Green, Missouri): total exchange capacity = 18.29 meq/100 g, pH = 5.8, organic matter = 2.7%, nitrogen = 74 lb/acre, sulfur (as sulfate) = 36 lb/acre, phosphorus (P) = 131 lb/acre, calcium (Ca) = 2,973 lb/acre, magnesium (Mg) = 707 lb/acre, potassium (K) = 264 lb/acre, sodium (Na) = 80 lb/acre, boron (B) = 3.16 lb/acre, iron (Fe) = 693.6 lb/acre, manganese (Mn) = 118.4 lb/acre, copper (Cu) = 2.8 lb/acre, zinc (Zn) = 5.6 lb/acre; percent base saturations: Ca = 54.3, Mg = 16.10, K = 1.85, Na = 0.95, other bases = 5.80%, H = 21.00.

Row width: 30 inches

Tillage: conventional

Experimental design: A small-plot corn study was conducted to evaluate the effect of Vitazyme and seaweed exact, using two applications, alone and in combination, on the yield of corn. There were four replications, with each plot four rows x 30 feet (10.00689 acres/plot); the two middle rows were harvested for analysis. A randomized complete block design was used.

Research organization: Agricultural Research and

Location: Cedar Falls, Iowa

Planting rate: 38,000 seeds/acre

Planting depth: 1.5 inches

Planting conditions: good

Previous crop: soybeans

Treatment	In-furrow at planting	Foliar
1. Control	0	0
2. Vitazyme	13 oz/acre	13 oz/acre
3. Seaweed	2 quarts/acre	2 quarts/acre
4. Vitazyme + Seaweed	13 oz/acre + 2 quarts/acre	13 oz/acre + 2 quarts/acre

Fertilization: All areas received 90 lb/acre of urea ammonium nitrate (UAN) on June 9.

Vitazyme application: (1) 13 oz/acre in-furrow at planting (May 8, 2014) using 38 ml/gallon at 10 gallons/acre; (2) 13 oz/acre on the leaves and soil at V6 (July 2, 2014), using 26 ml/gallon at 15 gallons/acre

Seaweed application: obtained from Ocean Organics; (1) 2 quarts/acre applied in-furrow at planting (May 8, 2014), using 189 ml/gallon at 10 gallons/acre; (2) 2 quarts/acre applied foliar after V6 (July 2, 2014) using 126 ml/gallon at 15 gallons/acre

Weather for 2014: Growing conditions for the trial were very good, with temperatures below normal and precipitation consistent throughout the crop cycle, except for a few days in June and August.

Irrigation: overhead center-pivot irrigation on June 12 and 13, and on August 11 to 14, and 18

Harvest date: October 21, 2014, using a Massey-Ferguson plot combine

Plant population results: Populations for the four treatments were very similar, without significant differences.


Treatment	Population*	Population change
	plants/acre	plants/acre
1. Control	34,065 a	—
2. Vitazyme	35,140 a	1,075 (+3%)
3. Seaweed	34,489 a	424 (+1%)
4. Vitazyme + Seaweed	34,111 a	46 (0%)
LSD (P = 0.05)	2,388	
CV	4.24%	
Treatment F	0.462	
Treatment probability	0.716	

*Means followed by the same letter are not significantly different at P = 0.05, according to the Student-Newman-Keuls Test.

Grain moisture results:

Treatment	Grain moisture*	Moisture change
	%	%
1. Control	17.66 a	—
2. Vitazyme	17.05 a	(-) 0.61
3. Seaweed	17.41 a	(-) 0.25
4. Vitazyme + Seaweed	17.14 a	(-) 0.52
LSD (P = 0.05)	0.78	
CV	2.82%	
Treatment F	1.276	
Treatment probability	0.340	


*Means followed by the same letter are not significantly different at P = 0.05, according to the Student-Newman-Keuls Test.



All three applications reduced grain moisture at harvest a small amount, with Vitazyme alone reducing moisture by 0.61 percentage-point.

Reduction in grain moisture with Vitazyme: 0.61 %-point

Grain yield results: The two center rows of each plot were harvested.



The three treatments all increased yield above the control, though not significantly, due to experimental error. Even so, Vitazyme increased yield by 5.5%.

Treatment	Corn yield*	Yield change
	bu/acre	bu/acre
1. Control	158.03 a	—
2. Vitazyme	166.70 a	8.67 (+5.5%)
3. Seaweed	165.78 a	7.75 (+4.9%)
4. Vitazyme + Seaweed	164.40 a	6.37 (+4.0%)
LSD (P = 0.05)	10.03	
CV	3.83%	
Treatment F	1.558	
Treatment probability	0.266	

*Means followed by the same letter are not significantly different at P = 0.05, according to the Student-Newman-Keuls Test.

Increase in corn yield with Vitazyme: 5.5%

Conclusions: A replicated corn study in east-central Iowa revealed that Vitazyme increased corn yield by 5.5%, followed closely by seaweed at 4.4%, and the combined products at 4.0%. Those increases were consistent, but no synergism was displayed between the two materials. Grain moisture was reduced by 0.61 percentage-point with Vitazyme, which gave the greatest reduction for the three treatments compared with the control. While these increases for the three treatments were not significant, they were consistently greater than the control so can be assumed to be real.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2014 Crop Results

Vitazyme on Corn

Researcher: unknown

Research organization: Kernel Company, LLC, Ukraine

Location: Man'kivs'ky District, Cherkasy Region, Viktorivka Village, Ukraine

Variety: DK 440

Planting rate: 75,000/ha

Planting date: April 8, 2014

Previous crop: soybeans

Soil type: Chernozem, with 3.7% organic matter

Seedbed preparation: disk-plowing to 6-8 cm, plowing to 22-24 cm, harrowing, two cultivations to 5 - 6 cm.

Experimental design: A corn field was divided into a Vitazyme treated area and an adjoining control area, with the objective of evaluating the effect of this product on soybean yield. All other treatments — herbicides and fertilizers — were identical over the entire area.

1. Control

2. Vitazyme


Fertilization: 90 kg/ha of nitrogen broadcast and incorporated before planting, and 10-26-26 kg/ha of N-P₂O₅-K₂O applied in-furrow at planting.

Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 5 to 6 - trifoliate stage, on May 3

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	7.91	—
Vitazyme	8.91	1.01 (+13%)

**Increase in corn yield
with Vitazyme: 13%**



Income results: The gross income minus expenses for the two treatments revealed the Vitazyme improved net income by 2,090 UAH/ha (\$132.72/ha at 1UAH = 0.0635 USD).

Increase in income with Vitazyme: 2,090 UAH/ha

Conclusions: This corn trial in Ukraine, using a single 1 liter/ha Vitazyme application at the 5 to 6-leaf stage, revealed a substantial 13% yield increase, which netted the farmer an extra 2,090 UAH/ha (\$32/72/ha). This result shows as have many other corn trials in Ukraine over the past years, the great efficacy of the Vitazyme program to boost yields and profits.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme and Fish on Corn

Responses at Four Nitrogen Levels

Researcher: Sara Berg and Ron Gelderman, Ph.D.

Research institution: Department of Agronomy and Soils, South Dakota State University, Brookings, South Dakota

Location: South Dakota State University Research Farm, Aurora, South Dakota

Variety: Dekalb 45-51

Soil type: Brandt silty clay loam

Soil test results: tested April 30, 2013; organic matter = 4.4%, NO₃-N = 32 lb/acre (to 24 inches), P (Olsen) = 20 lb/acre, K = 292 lb/acre, Zn = 2.56 lb/acre, S = 50 lb/acre (to 24 inches), pH = 5.5, salts (1:1) = 0.2 mmhos/cm

Planting date: May 15, 2013

Planting rate: 32,000 seeds/acre

Row spacing: 30 inches

Experimental design: A small plot (10 x 45 feet, or 0.01033 acre) design with four replications was used to evaluate the effectiveness of Vitazyme and liquid fish in enhancing corn yield, plant population, and grain nutrient content as influenced by nitrogen rate. The fish was applied alone and with Vitazyme at the high nitrogen rate only. Statistics were calculated for Treatments 1 to 8, and also for Treatments 7 to 10.

Treatment	Nitrogen rate	Vitazyme	Fish
	% of optimal		
1. Control, 0% N	0	0	0
2. Vitazyme, 0% N	0	X	0
3. Control, 50% N	50	0	0
4. Vitazyme, 50% N	50	X	0
5. Control, 75% N	75	0	0
6. Vitazyme, 75% N	75	X	0
7. Control, 100% N	100	0	0
8. Vitazyme, 100% N	100	X	0
9. Fish, 100% N	100	0	X
10. Fish + Vitazyme, 100% N	100	X	X

Fertilization: Nitrogen fertilizer (NH₄NO₃) was applied broadcast to the soil surface on May 15, just after planting. 100% N = 122 lb/acre of N; 75% N = 92 lb/acre of N; 50% N = 61 lb/acre of N.

Vitazyme application: (1) 13 oz/acre (1 liter/ha) in-furrow at planting, with 3 gallons of water per acre along with 10-34-0% N-P₂O₅-K₂O fertilizer; (2) 13 oz/acre (1 liter/h)a sprayed with a hooded plot sprayer on the leaves and soil at V8, on July 10, 2013


Fish application: (1) 2 gal/acre in a 5 gal/acre volume, with Vitazyme and 10-34-0% N-P₂O₅-K₂O, in-furrow at planting; (2) 2 gal/acre sprayed on the leaves and soil with a hooded plot sprayer at V8, on July 10, 2013; (3) 2 gal/acre on the leaves, hand-sprayed with a backpack sprayer at silking

Harvest date: October 25, 2013


Chlorophyll results: Chlorophyll readings were made with a Minolta SPAD meter at V10 (July 17, 2013), and again at R1 (August 1, 2013). No consistent results were noted, although the highest overall value recorded was for 100% N + Vitazyme at R1 (54.6), which was 1.2 points higher than the respective control treatment.

Population results: There was some benefit to plant stand shown with Vitazyme at three out of the four nitrogen levels.

Plant Population and Nitrogen Rate



Plant Population With Vitazyme and Fish




In every case except at 0% N, the Vitazyme and fish treatments exceeded the control treatments. Fish alone and Vitazyme + fish significantly exceeded the control in plants/acre.

Grain moisture results: There were no clear relationships between Vitazyme, fish, and nitrogen rates. At harvest moisture was around 16.0 to 17.0%


Grain test weight results: Weight per bushel of grain varied little across all of the treatments, ranging from 53.4 to 55.9 lb/bu for the eight nitrogen rate treatments. The Vitazyme + Fish treatment, however produced the heaviest grain at 56.4 lb/bu.

Grain yield results:

Grain Yield and Nitrogen Rate



Grain Yield With Vitazyme and Fish



*Significantly greater than the control treatment at the same nitrogen level at P = 0.05, according to the Statistical Analysis System, GLM.

*Significantly greater than the control treatment at the same nitrogen level at P = 0.05, according to the Statistical Analysis System, GLM.

Increase in Grain Yield With Vitazyme

0% N	8.7%
50% N	1.3%
75% N	4.4%
100% N	3.5%

Grain yield increases were consistent across all nitrogen levels, but the largest percentage increase (8.7%) was with no fertilizer nitrogen added.


Increase in Grain Yield With Vitazyme and Fish - 100% N

Vitazyme only	3.5%
Fish only	3.2%
Vitazyme + Fish	4.5%

Both Vitazyme and fish alone significantly improved grain yield, but the combined Vitazyme and fish produced the highest yield — 4.5% higher than the 100% control — which was the highest yield of all ten treatments (164.2 bu/acre).


Grain quality results: Analyses of oil, protein, and starch were performed at the South Dakota State University diagnostic laboratory. Oil varied from 2.94 to 3.29%, protein from 5.26 to 7.53%, and starch from 57.74 to 58.87%. There was no effect of Vitazyme and fish on these contents although nitrogen levels increased the protein for the control and Vitazyme treatments. See the following graph.

Protein of Grain and Nitrogen Level




Analyses of grain potassium, copper, iron, zinc, manganese, phosphorus, sulfur, magnesium, and calcium revealed no clear patterns of effects for Vitazyme and fish on these nutrients, although fish tended to increase potassium and phosphorus. Nitrogen increased some nutrient levels and reduced others. Values of both the control and Vitazyme treatments are averaged for each N level.


Grain K and N Level




Grain Cu and N Level




Grain Fe and N Level




Grain Zn and N Level




Grain Mn and N Level




Grain P and N Level




Grain S and N Level



Grain Mg and N Level



Grain Ca and N Level



Conclusions: A small plot corn study in east central South Dakota revealed that Vitazyme and fish — Vitazyme applied at planting and at V8, and fish applied at planting, V8, and silking — significantly increased corn yield at all nitrogen fertilizer levels, by up to 8.7% at the 0% N level. The Vitazyme plus fish treatment at 100% N gave the highest yield of all ten treatments (164.2 bu/acre). Vitazyme increased plant population at the three highest nitrogen levels, and fish and fish plus Vitazyme also increased population at 100% N. Grain quality parameters were affected by nitrogen levels to some degree, the contents of protein, copper, iron, manganese, and sulfur increasing and the contents of potassium, zinc, phosphorus, and magnesium decreasing with increasing nitrogen fertilizer. This study reveals the usefulness of Vitazyme and fish, and especially the two combined, to increase corn yields under rainfed conditions in eastern South Dakota.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Farmer: Ramiro Carlos Fonseen

Location: El Ahuaje de Abajo Farm, San Nicolas, Cocula, Jalisco, Mexico

Variety: N 432 HJ (Eagle)

Planting date: May 8, 2012

Experimental design: Corn to be used for leaves to wrap tamales was treated with Vitazyme on the seeds, and later on the leaves and soil, to one portion of a field, and compared with an untreated control to evaluate leaf production. The treated area was 1 ha.

1. Control

2. Vitazyme

Fertilization: unknown

Vitazyme application: (1) Seed treatment of 50 ml in 450 ml of water, to treat 55,000 seeds (a 20 kg bag planted 1 ha); (2) soil and foliar treatment of 1 liter/ha (13 oz/acre) at an unknown date.

Leaf yield result: Actual yield numbers for the two plots were not available, but the data reported an 8% increase in leaves with Vitazyme versus the control.

Increase in yield with Vitazyme: 8%

Conclusion: This Mexican corn study, utilizing two Vitazyme applications — a seed and a soil/foliar treatment — showed that this product significantly enhanced leaf production of the corn plants.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Location: Camino a la Coronilla Farm, Ameca, Jalisco

Planting date: May 8, 2012

Experimental design: A 1 ha plot was selected from a corn field to treat with Vitazyme, on both the seeds and leaves, to determine the effectiveness of the product to increase yield.

Farmer: Eng. Jose Guadalupe Orozco Flores

Variety: Pioneer 3055

Planting rate: 80,000 seeds/ha

1. Control

2. Vitazyme


Fertilization: unknown

Vitazyme application: (1) A seed treatment of 50 ml in 450 ml of water, applied with Regent; (2) a soil and foliar spray treatment of 1.0 liter/ha at an unknown date

Yield results:

Treatment	Grain yield tons/ha	Yield change tons/ha
Control	11.1	—
Vitazyme	13.0	1.9 (+17%)

**Increase in grain yield with
Vitazyme: 17%**



Conclusions: A corn study with Vitazyme in Mexico proved that the product increased grain yield substantially, by 17%, over the untreated control, showing the great effectiveness of this material to stimulate corn yield.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Tim Heikens and Leonard Jansen

Location: Lake Park, Iowa

Row spacing: 30 inches

Planting date: May 11, 2013

Experimental design: A 30-acre uniform corn field was sprayed with Vitazyme in alternating 90-foot sprayer strips through the field after the corn was growing. The purpose of the study was to determine if a single application could improve grain yield.

Farmer: Leonard Jansen

Variety: Golden Harvest H7891

Plant population: 33,000 seeds/acre

Soil type: silty clay loam (Mollisol)

1. Control

Fertilization: Total nutrients applied were 140-100-120 lb/acre of N-P₂O₅-K₂O.

Vitazyme application: 20 oz/acre (1.5 liters/ha) sprayed on the leaves and soil on July 6 at the 10 to 11 leaf stage

Weed control: 1.5 pints/acre Steadfast on May 9; 1 quart/acre Roundup on June 17

Weather during the season: a wet spring and a dry summer and fall

Harvest date: October 24, 2013


Yield results: A weigh wagon was used to quantify yields.

2. Vitazyme

Treatment	Yield bu/acre	Yield change bu/acre
Control	181.19	—
Vitazyme	184.16	2.97 (+2%)

Grain moisture results: the Vitazyme treated corn had 17% moisture and the control corn had 18% moisture.

**Reduction in grain moisture
with Vitazyme: 1.0%**



Conclusions: This corn trial in northwestern Iowa revealed that Vitazyme, applied at the 10 to 11-leaf stage at 20 oz/acre, produced about a 3 bu/acre yield increase. Grain moisture was 1.0% less. Had the product been applied at planting as well the yield improvement would likely have been much greater.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: unknown
Nong, Viet Nam

Farmer: Linh Thi Ngan
Variety: P4199

Location: Dak Will, Cu Jut District, Dak
Planting date: August 10, 2013

Planting rate: 50,000 plants/ha

Experimental design: A corn field was divided into a Vitazyme treated and control area to determine the effects of this product on corn growth and yield.

1. Control

2. Vitazyme

Fertilization: unknown

Vitazyme application: (1) seed treatment with a 5% solution before planting; (2) 2 liters/ha spray on the leaves and soil 15 days after planting; (3) 2 liters/ha spray on the leaves and soil 30 days after planting.

Germination results: Both treatments had over 95% germination.

Harvest date: December 12, 2013, 120 days after planting

Harvest and Yield results:

Treatment	Plant height	Ear height	Drought resistance	Weight of 1,000 seeds	Seed color	Yield
	cm	cm		grams		tonnes/ha
Control	185	100	Fair	310	Light yellow	8.0
Vitazyme	192 (+4%)	110 (+10%)	Good	315 (+2%)	Dark yellow	9.0 (+13%)

Increase in plant height: 4%

Increase in ear height: 10%

Increase in 1,000 seed weight: 2%

Increase in yield: 13%

Income results: The Vitazyme treated corn sold for 4,000 VND/kg, versus 3,800 VND/kg for the control corn.

Costs with Vitazyme: 1,420,000 VND/ha

Increased income with Vitazyme: 4,180,000 VND/ha

Return On Investment with Vitazyme: 2.94 : 1

Conclusions: A corn study in Viet Nam proved that a Vitazyme seed treatment, plus two soil and foliar applications, produced an excellent 13% yield increase. Drought tolerance was improved, as was seed weight (2%), and plant height (4%). Income was substantially increased, giving a Return On Investment of 2.94 : 1. This program is an excellent one for corn production in Viet Nam.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Location: La Mesita Farm, Cocula, Jalisco, Mexico

Planting date: 90,000 seeds/ha

Experimental design: A 1.0 ha area of an irrigated corn field was treated with Vitazyme on the seeds, and later on the leaves and soil, in order to evaluate the product's ability to increase grain yield.

1. Control

Fertilization: unknown

Vitazyme application: (1) Seeds were treated with a 10% solution; (2) leaves and soil were sprayed with 1 liter/ha (13 oz/acre) during growth at an unspecified time.

Yield results:

Treatment	Grain yield tons/ha	Yield change tons/ha
Control	12.8	—
Vitazyme	15.3	2.5 (+20%)


**Increase in grain yield with
Vitazyme: 20%**

Farmer: Eng. Juan Carlos Flores Orozco

Variety: Pioneer 3055

Irrigation: sprinkler

2. Vitazyme



Conclusions: A corn study under irrigation, using a seed treatment and an additional foliar and soil spray, revealed that Vitazyme improved grain yield by 20% above the untreated control.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme and Kelpek on Corn

Researcher: V.V. Plotnikov Research organization: Scientific, Innovation, and Technology Center of the Institute of Forages and Agriculture of Podillya NAAS Location: National Academy of Agricultural Sciences, Ukraine
Variety: DKS 2960, FAO 250 Soil type: ash gray soil (humus = 2.2%, hydrolyzed-N = 8.4 mg/100 g of soil, P = 15.8 mg/100 kg of soil, exchangeable K = 12.4 mg/100 g of soil, pH = 5.5)
Previous crop: winter wheat Planting date: May 10, 2013
Soil preparation: disking, plowing, harrowing, cultivating Seeding rate: 80,000 seeds/ha
Experimental design: A small plot experiment with four replications was prepared to evaluate the effects of Vitazyme and Kelpek on the yield and growth of corn.

1. Control

2. Vitazyme


3. Kelpek

Fertilization: Before planting, 100-60-60 kg/ha of N-P₂O₅-K₂O were applied to the active test area.

Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 7 to 8-leaf stage (June 11, 2013)

Kelpek application: 2 liters/ha sprayed on the leaves and soil at the 7 to 8-leaf stage (June 11, 2013)

Yield results:



Increase in yield	
Vitazyme	15%
Kelpek	8%


Both Vitazyme and Kelpek substantially increased corn yield, but Vitazyme produced nearly double the increase of Kelpek (15%)

Income results:

Increase in net income	
Vitazyme	1,690 UAH/ha
Kelpek	780 UAH/ha

Vitazyme clearly produced the greater income of the two products, more than doubling that achieved by Kelpek.

Leaf area results:



Increase in leaf area
Vitazyme 35%
Kelpek 19%

Vitazyme nearly doubled the leaf area of corn versus Kelpek.

Conclusions: The authors of the report stated,

1. The treatment of corn hybrid DKS 2960 at the 7-8 leaf stage with Vitazyme, at 1 L/ha, gave a corn grain yield increase of 106 tonnes/ha, or 15%. When applying the competitive chemical Kelpek at a rate of 1 L/ha at the 7-8 leaf stage, the increase of corn grain yield was 0.9 tonne/ha, or 8%, and was 0.7 tonne/ha less than with Vitazyme.
2. With Vitazyme at 1 L/ha, the profit was 1,690 UAH/ha, which is 910 UAH/ha more than when applying Kelpek at 2 L/ha.
3. Vitazyme application at 1 L/ha gave an increase in corn leaf area of 1.1 m²/m² of the plot. Applying Kelpek at 2 L/ha gave a corn leaf area increase of 0.6 m²/m² of the plot, which is 0.5 m²/m² less than when applying Vitazyme.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Location: San Juan Acozac, Puebla, Mexico

Experimental design: A corn field for fodder was divided into Vitazyme and control areas of 4,000 m² (0.4 ha) each, to evaluate the effect of the product on corn yield.

Farmer: Odilon Ramos

Variety: unknown

1. Control

2. Vitazyme


Fertilization: unknown

Vitazyme application: 1 liter/ha (13 oz/acre) two times (timing unknown)

Yield results: Two cuttings were weighed for each treatment.

Treatment	Forage yield		Yield change
	tons	tons/ha	tons/ha
Control	1.9	4.8	—
Vitazyme	4.0	11.9	7.1 (+148%)

**Increase in forage yield with
Vitazyme: 148%**



Conclusions: A forage corn study in Mexico revealed that Vitazyme applied twice during the growing season produced a remarkable 111% increase in yield. This program is highly effective for corn growers in Mexico.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Farmer: Eng. Salvador Sanchez Rica

Location: El Chivero Field, El Llano Farm, San Martin Hidalgo, Jalisco, Mexico

Variety: Cimarron

Experimental design: A corn field had 1 ha. selected to be treated with Vitazyme, applied to the seeds and later as a foliar spray, to evaluate effects on yield compared to an adjacent untreated area.

1. Control

2. Vitazyme


Fertilization: unknown

Vitazyme application: (1) Seeds were treated with 100 ml/20 kg of seed; (2) a spray of 1 liter/ha (13 oz/acre) was made to the leaves and soil sometime during growth.

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	7.30	—
Vitazyme	8.21	0.91 (+12%)

**Increase in yield with
Vitazyme: 12%**



Conclusions: This Mexican corn study revealed that a Vitazyme seed treatment and single foliar treatment increased grain yield by 12%, showing the program to be highly valuable for corn production in Mexico.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme and Fish on Corn

Researcher: Bertel Schou, Ph.D.

and Educational Services), Cedar Falls, Iowa

Variety: Pioneer P0453HR (Roundup Ready)

Planting rate: 38,000 seeds/acre

Planting date: May 17, 2013

Tillage: conventional (field cultivating and harrowing)

Soil test values: pH = 6.9, organic matter = 2.7%, cation exchange capacity = 20.0 meq/100 g, N = 74 lb/acre, SO₄-S = 16 lb/acre, P₂O₅ = 246 lb/acre, Ca = 5,743 lb/acre, Mg = 807 lb/acre, K₂O = 302 lb/acre, Na = 94 lb/acre, B = 1.45 lb/acre, Fe = 539 lb/acre, Mn = 143.7 lb/acre, Cu = 2.3 lb/acre, Zn = 11.8 lb/acre; percent base saturations: Ca = 71.6%, Mg = 16.8%, K = 1.9%, Na = 1.0%, other bases = 4.6%, H = 4.0%.

Experimental design: A corn study having individual plots 15 x 30 feet (450 ft², or 0.0103 acre) was set up in a randomized complete block design, with six replicates. The purpose of the study was to determine the effects of Vitazyme and fish applied separately, and the two applied together in-furrow and foliar, on crop yield, moisture, and plant population.

Research organization: ACRES (Agricultural Research

Location: Cedar Falls, Iowa BBCH scale: BCOR

Maturity: 104 days

Row spacing: 30 inches

Previous crop: soybeans

Planting Depth: 1.5 inches

Slope of plot: 5%

Soil type: Kenyon loam

Treatment	At planting	At V6R1	At R4
1. Control	0	0	0
2. Vitazyme	13 oz/acre	13 oz/acre	0
3. Fish	2 gal/acre	2 gal/acre	2 gal/acre
4. Vitazyme + Fish	13 oz + 2 gal (Trt. 2 + 3)	13 oz + 2 gal (Trt. 2 + 3)	2 gal/acre (Fish only)

Fertilization: In the fall of 2012, 100 lb/acre of 18-46-0 (% N-P₂O₅-K₂O) and 100 lb/acre of 0-0-60. At 14 inches corn height, 100 lb/acre of N was applied as UAN (urea ammonium nitrate).

Vitazyme application: At planting (for Treatments 2 and 4), 13 oz/acre (1 liter/ha) in-row in a 10 gallon/acre solution; at V6 (Treatments 2 and 4), 13 oz/acre (1 liter/ha) sprayed on June 20 at 15 gallons/acre.

Fish application: Emulsified fish was applied at 2 gallons/acre in-row at planting (Treatments 3 and 4), and at 2 gallons/acre by foliar spray at V6 (Treatments 3 and 4) on June 20, as well as at V10 (Treatments 3 and 4) on August 26. The fish was mixed with water to give a 20 gal/acre output.


Weed control: Harness Xtra preplant (1.2 quarts/acre); Roundup postemergence at the recommended rate

Harvest date: October 25, 2013

Yield results: Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Kuels Test.

Treatment	Yield ¹ bu/acre	Yield change bu/acre
1. Control	240.3 b	—
2. Vitazyme	248.1 ab	7.8 (+3.2%)
3. Fish	249.3 a	9.0 (+3.7%)
4. Vitazyme + Fish	247.9 ab	7.6 (+3.2%)
LSD (P = 0.05)	8.0	
Standard deviation	6.5	
CV	2.6%	
Replicate F	1.73	
Treatment F	2.40	

¹Adjusted to 15.5% moisture.



Increase in corn yield with Vitazyme: 3.2%

Increase in corn yield with Fish: 3.7%


Increase in corn yield with Vitazyme + Fish: 3.2%

Although only the fish treatment statistically exceeded the control, the three Vitazyme and fish treatments were equal, differing by 0.5% from each other. Vitazyme increased yield at the 6% confidence level, but the combined Vitazyme and fish did not enhance the yield above either product alone.

Plant population results: All final population values were nearly equal (36,459 to 36,776 plants/acre).

Grain moisture results:

Treatment	Grain moisture %	Moisture change %
1. Control	18.92 b	—
2. Vitazyme	19.22 ab	+0.30
3. Fish	19.09 ab	+0.17
4. Vitazyme + Fish	19.73 a	+0.81
LSD (P = 0.05)	0.62	
Standard deviation	0.50	
CV	2.6%	
Replicate F	1.73	
Treatment F	2.92	



Only the combined Vitazyme + Fish treatment exceeded the control in grain moisture at P = 0.05. The other treatments were very close to each other.

Conclusions: A replicated corn trial in east-central Iowa revealed that both Vitazyme and liquid fish, and the products combined, increased the yield of corn grain significantly — Vitazyme alone and the two products together at P = 0.06, and fish alone at P = 0.05 — with the combined products not showing a synergistic effect. These yield increases were 7.8, 9.0, and 7.6 bu/acre for the Vitazyme, fish, and combined treatments, respectively. Plant population was not affected by either material, but grain moisture was greater than the control for the combined products by 0.81 percentage point at P = 0.05.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

Researcher: Juan Carlos Diaz, Ph.D.

Farmer: Eng. Jose Isabel Hernandez Soto

Location: El Monte Farm, Villa Corona, Jalisco, Mexico

Variety: Cayman

Planting date: May 8, 2012

Experimental design: A corn field was divided into Vitazyme treated and control areas to determine the effect of a seed and foliar treatment on corn yield. The treated area was 1 ha.

1. Control

2. Vitazyme


Fertilization: unknown

Vitazyme application: (1) Seed treatment, at 100 ml/20 kg of seed; (2) soil and foliar spray at 1 liter/ha (13 oz/acre) at an undetermined time

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	8.6	—
Vitazyme	10.5	1.9 (+22%)

**Increase in yield with
Vitazyme: 22%**



Conclusions: Applying a seed treatment and foliar spray to this corn study resulted in an excellent 22% yield increase.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

A Nitrogen Rate Study

Researcher: Timothy Veldkamp

Research Organization: SGS Ag Research

Location: Aurora, South Dakota

Variety: 4055 Roundup Ready

BCCH Scale: BCOR

Planting Date: June 14, 2013

Planting rate: 32,000 seeds/acre

Planting depth: 1.5 inches

Row spacing: 30 inches

Tillage: conventional

Previous crop: soybeans

Soil type: Strayhoss Loam (pH = 6.5, organic matter = 4.5%, cation exchange capacity = 25 meq/100 g of soil, P = 14 lb/acre, K = 174 lb/acre, Zn = 1.22 lb/acre)

Experimental design: A small plot soybean study was designed to evaluate the effect of Vitazyme, at two nitrogen levels, on the yield of corn. The plots were 10 x 40 feet (0.009183 acre), and arranged in a randomized complete block design, with four replicates.

1. Control, 50% N 2. Vitazyme, 50% N 3. Control, 100% N 4. Vitazyme, 100% N

Fertilization: 50% N plots received 65 lb/acre of N; 100% N plots received 130 lb/acre of N

Vitazyme application: (1) 13 oz/acre (1 liter/ha) in the seed row at planting; (2) 13 oz/acre (1 liter/ha) on the leaves and soil at V8.

Weed control: 1.8 quarts/acre of Harness Xtra + 24 oz/acre of Roundup PowerMax on July 3; 24 oz/acre of Roundup PowerMax on August 8

Vigor results: Plant vigor was evaluated at the V2 stage on July 8, 2013. Values ranged from 5.8 to 6.3, and were not significantly different.

Harvest date: November 8, 2013


Yield results: The middle two rows of each plot were combined and weighed, and the moisture content was determined using an Almaco Grain Gauge.

Treatment	Grain yield ¹ bu/acre	Yield change ² bu/acre
Control, 50% N	159.0 b	—
Vitazyme, 50% N	165.3 a	6.3 (+4%)
Control, 100% N	167.4 a	—
Vitazyme, 100% N	170.3 a	2.9 (+2%)
LSD (P = 0.10)	5.0	
CV	2.35%	
Replicate F	13.880	
Treatment F	6.138	

¹Means followed by the same letter are not significantly different at P=0.10, according to the Student-Newman-Kuels Test.

²The Vitazyme means are compared with the control means at the same N level.

Grain Yield, 50% N



Increase in yield at 50% N: 4%

Increase in yield at 100% N: 2%

Vitazyme significantly increased grain yield at 50% N, boosting it by 6.3 bu/acre, while the 100% N application revealed that Vitazyme increased the yield by 2.9 bu/acre over the control, but not significantly. Vitazyme grain yield at 50% N was statistically the same as was the untreated grain yield at 100% N.


Test weight results: All values varied within a narrow range — 51.15 to 51.93 lb/bu — and none of the differences were significant. However, Vitazyme at both N levels slightly raised bushel weight.

**Increase in test weight
with Vitazyme**

At 50% N 0.78 lb/bu

At 100% N 0.25 lb/bu

Grain Yield, 100% N



Grain moisture results: There were no significant differences in grain moisture among the four treatments.

Conclusions: This corn trial with Vitazyme at two nitrogen levels, near Brookings, South Dakota, revealed that Vitazyme significantly improved corn yield at the 50% N level. This yield was only 2.1 bu/acre less than the 100% N treatment without Vitazyme, and not significantly different from it. This reveals the improved nitrogen efficiency often noted with the use of this product. The 100% N level gave a 2% (2.9 bu/acre) yield increase for Vitazyme; the two were statistically the same. Grain moisture was not affected by the treatments, but there appeared to be a slight improvement in grain bushel weight with Vitazyme. These results show the excellent value of this program for corn growers in the northern Corn Belt.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2013 Crop Results

Vitazyme on Corn

With Quantum Products at Two Nitrogen Rates

Researcher: Bertel Schou, Ph.D.
and Educational Services), Cedar Falls, Iowa

Variety: Pioneer P9675 (Roundup Ready)

Planting rate: 38,000 seeds/acre

Tillage: field cultivated and harrowed

Soil type: Maxfield silty clay loam (organic matter = 4.4%, pH = 6.3, cation exchange capacity = 17 meq/100 g of soil, drainage = excellent, fertility = excellent)

Experimental design: A small plot corn experiment was established in east-central Iowa to evaluate the effect of Vitazyme and two Quantum products on the yield of corn, at two nitrogen levels. Six replications were used, with a randomized complete block design. Individual plots were 15 x 30 feet, or 450 ft² per plot (0.01033 acre). This experiment had been put in earlier but was mistakenly fertilized incorrectly, so was redone and planted later with a short-season variety, so yielded less than the original study would have.

Research organization: ACRES (Agricultural Research

Location: Cedar Falls, Iowa

BBCH scale: BCOR

Planting Depth: 1.5 inches

Maturity: 96 days

Row spacing: 30 inches

Slope of plot: 4%

Planting date: June 20, 2013

Previous crop: corn

Treatment	Nitrogen	Vitazyme	Quantum Light	Quantum VSC
1. Control	50%	0	0	0
2. Vitazyme	50%	In-furrow,Foliar	0	0
3. Quantum	50%	0	In-furrow, Foliar	In-furrow, Foliar
4. Vitazyme + Quantum	50%	In-furrow, Foliar	In-furrow, Foliar	In-furrow, Foliar
5. Control	100%	0	0	0
6. Vitazyme	100%	In-furrow, Foliar	0	0
7. Quantum	100%	0	In-furrow, Foliar	In-furrow, Foliar
8. Vitazyme + Quantum	100%	In-furrow, Foliar	In-furrow, Foliar	In-furrow, Foliar

Fertilization: 18-46-0% N-P₂O₅-K₂O applied per acre the fall of 2012; 50% N, 80 lb/acre urea-NH₄NO₃; 100% N, 160 lb/acre urea NH₄NO₃, all sidedressed

Vitazyme application: 13 oz/acre (1 liter/ha) in-row at planting; 13 oz/acre (1 liter/ha) on the leaves and soil at V8, August 9, 2013

Quantum application: Both Quantum products are produced by Applied and Experimental Biology, Jacksonville, Florida. Quantum Light is a proprietary blend of beneficial cultures and photosynthetic microbes that increase plant photosynthesis and growth. Quantum VSC contains photosynthetic microbes, hypercellulose, symbiotic microbes, and humic substances. These were applied at 1 gallon/acre each for both in-furrow and foliar applications, at the same times as Vitazyme. The two were applied together.

Weed control: post-emergence herbicides

Weather during the growing season: Temperatures were about average after a cool early season; spring was wet, but July and August were dry.


Harvest date: October 28, 2013

Yield results: The two center rows of each plot were harvested for yield results. Weights are corrected to 15.5% moisture.


Treatment	Grain yield ¹ bu/acre	Yield change ² bu/acre
----- 50% Nitrogen -----		
1. Control	101.9 d	—
2. Vitazyme	116.4 c	14.5 (+14%)
3. Quantum	120.0 c	18.1 (+18%)
4. Vitazyme + Quantum	125.3 bc	23.4 (+23%)
----- 100% Nitrogen -----		
1. Control	116.4 c	—
2. Vitazyme	131.8 ab	15.4 (+13%)
3. Quantum	134.2 ab	17.8 (+15%)
4. Vitazyme + Quantum	136.7 a	20.3 (+17%)
LSD (P = 0.05)	10.4	
CV (P = 0.05)	7.21%	
Replicate F	4.051	
Treatment F	10.244	

¹Means followed by the same letter are not significantly different at P = 0.05 according to the Student-Newman-Kuels Test.
²Comparisons are made within the same fertility level.

Yield at 50% Nitrogen



Yield at 100% Nitrogen



Vitazyme consistently boosted yield by 13 to 14% above the control at both nitrogen levels, while the Quantum products improved the yield by 15 to 18%. The two products together further boosted corn yield, to 17 to 23% above the control. In every cases the greatest increases were at the 50% nitrogen level. There appears to be a synergism between Vitazyme and Quantum materials, with additional boosts in yield of 2% at the 100% N level, and 5% at the 50% N level.

	50% N	100% N
Vitazyme	14%	13%
Quantum Only	18%	15%
Vitazyme + Quantum	23%	17%

Grain moisture results: Grain moisture was determined by a device on the plot combine.


Treatment	Grain moisture ¹	Change ²
	%	%e
----- 50% Nitrogen -----		
1. Control	25.00 a	—
2. Vitazyme	23.93 ab	-1.07
3. Quantum	23.38 ab	-1.62
4. Vitazyme + Quantum	23.30 b	-1.70
----- 100% Nitrogen -----		
1. Control	23.89 ab	—
2. Vitazyme	24.16 ab	+0.27
3. Quantum	24.35 ab	+0.46
4. Vitazyme + Quantum	24.02 ab	+0.13
LSD (P = 0.05)	1.67	
CV	5.90%	
Replicate F	0.940	
Treatment F	0.869	

¹Means followed by the same letter are not significantly different at P = 0.05 according to the Student-Newman-Kuels Test.
²Comparisons are made within the same fertility level.

There was only one significant grain moisture change: a reduction of 1.70 percentage points with Vitazyme + Quantum at 50% N. Both Vitazyme and Quantum and Quantum alone also reduced grain moisture, but not significantly at 50% N. On the other hand, all treatments at 100% N slightly increased grain moisture.

Reduction in grain moisture with Vitazyme + Quantum at 50% N: 1.70%-point

Change in Grain Moisture



Plant population results: All treatments had statistically the same population (23,980 to 22,092 plants/acre) except the control at 50% N, and Quantum products at 50% N, and Quantum products at 50% N, which had significantly lower populations. The good yield response of the Quantum treatment showed that these slightly reduced populations had little effect on the resultant yield.

Conclusions: A small plot corn study in east-central Iowa revealed that a late-planted Roundup Ready variety responded well to both two nitrogen levels, and also Vitazyme and Quantum products, alone or in combination. Vitazyme boosted the grain yield by 14% at 50% N, and by 13% at 100% N; Quantum Light + Quantum VSC increased the yields by 18% and 15% at 50 and 100% N, respectively. The products applied together in-row and foliar brought a synergistic improvement in yield above either product alone, the increases being 23% at 50% N and 17% at 100% N. Grain moisture was significantly reduced by the combined products at the 50% N level, but this reduction was not achieved at the 100% N level. Both products, and especially the products applied together, showed fine grain yield responses in this trial, and should be evaluated further as to their potential as excellent synergists for farmers in the Corn Belt.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Researcher: James Anderson

Farmer: David Bergeson

Location: Dawson, Minnesota

Varieties: Pioneer 38A56, Garst 88B37

Planting date: April 25, 2012

Row width: 30 inches

Planting rate: 32,000 seeds/acre

Previous crop: soybeans

Tillage: conservation

Experimental design: A field was divided into plots of field length to determine the effects of Vitazyme on corn yield and test weight for various varieties.

1. Control

2. Vitazyme

Fertilization: (1) starter in-furrow at 8 gal/acre (a 20-14-12% N-P₂O₅-K₂O formulation with sulfur and zinc, dry mixed at 1.5 lb/gal of water); (2) 120 lb/acre of anhydrous NH₃ (82% N)

Vitazyme application: 13 oz/acre in the starter, on the seed row at planting

Weather: very dry all season

Yield and test weight results: A weigh wagon was used to weigh the corn for a measured area, from which yield calculations were made. Harvest date was September 29, 2012.

Variety: Pioneer 38A56



Treatment	Yield	Yield change	Test weight	Test weight change
	bu/acre	bu/acre	lb/bu	lb/bu
Control	146.5	—	57.9	—
Vitazyme	155.0	8.5 (+6%)	59.5	+1.6

Yield

Test Weight

**Increase in yield
with Vitazyme: 6%**

**Increase in test
weight with
Vitazyme: 1.6 lb/bu**





Pioneer 38A56 responded with a yield increase of 6% (8.5 bu/acre), and with heavier grain that weighed 1.6 lb/acre more with Vitazyme.

Variety: Garst 88B37

Treatment	Yield	Yield change	Test weight	Test weight change
	bu/acre	bu/acre	lb/bu	lb/bu
Control	143.0	—	51.9	—
Vitazyme	155.3	12.3 (+9%)	53.8	+1.9

Increase in yield with Vitazyme: 9%

Increase in test weight with Vitazyme: 1.9 lb/bu




Garst 88B37 produced excellent responses with Vitazyme, the yield increasing by 9% (12.3 bu/acre), and the test weight by 1.9 lb/bu.

Variety: Pioneer 36152

Treatment	Yield	Yield change
	bu/acre	bu/acre
Control	124.2	—
Vitazyme	129.2	5.0 (+4%)


Increase in yield with Vitazyme: 4%



Variety: Federal 5100

Treatment	Yield	Yield change
	bu/acre	bu/acre
Control	84.4	—
Vitazyme	105.0	20.6 (+24%)

Increase in yield with Vitazyme: 24%



The great increase in yield with Vitazyme was due in part to the fact that the harvest was postponed because of equipment problems. As a result, high winds blew down some of the corn, preferentially downing the untreated control corn because the stalks and ear shanks were weaker

Conclusions: Four comparisons of Vitazyme treated and untreated corn varieties in this west central Minnesota study produced excellent yield increases for Vitazyme: 6% for Pioneer 38A56, 9% for Garst 88B37, 4% for Pioneer 36152. and 24% for Federal 5100. The 24% increase for the Federal variety was due in part to better standability and stronger stalks and shanks of the great value in using Vitazyme to increase corn yields, grain weight, and profits in western Minnesota. Since only 13 oz/acre was used at planting, at a cost of about \$5.00/acre, then the smallest increase (5.0 bu/acre) netted an added \$32.50/acre — using \$7.50/bu of corn — while the highest increase (20.6 bu/acre) netted \$149.50/acre more with a single Vitazyme application. The recommended two applications would likely have given addition returns on investment.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Free Amino Acid Levels - University of Missouri

Researcher: Manjula Nathan, Ph. D. Location: University of Missouri, Columbia, Missouri
Variety: unknown Planting date: May 3, 2011 Row spacing: 30 inches
Seeding rate: 32,000 seeds/acre Soil type: unknown

Experimental design: A plot area of 12 rows — six rows for the control and six rows for the Vitazyme treatment (0.33 acre per treatment) — was selected; each row was 200 feet long. Corn was grown to determine yield responses (reported elsewhere), and also to evaluate the free amino acid levels in plant tissues.

1. Control

2. Vitazyme

Fertilization: station standard for corn

Vitazyme application: The seeds for Treatment 2 were treated with 100% Vitazyme to give a coating of 13 oz on 32,000 seeds, or an equivalent of 13 oz/acre. The control received only water on the seeds. On June 28 (V10 stage), an additional 13 oz/acre was applied to the leaves and soil of Treatment 2.

Free amino acid analysis: Samples of leaves below and opposite the ear were collected on July 14, during silking. These leaves were quite ragged due to the hail damage of a severe storm that struck on July 3, 11 days before. These composite samples from each treatment were frozen and later analyzed at a university laboratory to determine levels of free amino acids.

Amino acid*	Control	Vitazyme	Change
	µg/100 mg	µg/100 mg	µg/100 mg
Aspartic acid	23.32	20.04	-3.28
Threonine	21.72	20.33	-1.39
Serine	16.13	15.52	-0.61
Glutamic acid	29.55	28.15	-1.40
α-amino adipic acid	0.78	1.45	+0.67
Proline	9.49	9.73	+0.24
Gycine	6.88	5.84	-1.04
Alanine	35.26	32.81	-2.45
Valnine	14.06	13.40	-0.66
Methionine	0.61	0.46	-0.15
Cystine	11.07	11.43	+0.36
Isoleucine	10.21	15.39	+5.18
Leucine	15.28	16.64	+1.36
Tyrosine	11.14	11.83	-0.69
Phenylalanine	12.37	10.11	-2.26
β-amino isobutyric acid	5.80	4.66	-1.14
alpha-amino butyric acid	6.63	6.12	-0.51
Ornithine	3.29	7.93	+4.64
Lysine	14.89	13.80	-1.09
Histidine	3.26	3.10	-0.16
Arginine	0.72	3.20	+2.48
Net change			-1.90

*Only those free amino acids that were discovered in the analysis are listed here.

Note that the net effect of Vitazyme is to decrease the free amino acid level of the leaf tissue slightly (1.90 µg/100mg) This occurred in spite of the fact that large increases occurred for isoleucine, ornithine, and arginine. Most amino acids decreased with Vitazyme treatment, which should help explain the observation that treated plants resist fungal, bacterial, nematode, and other infections.

Conclusions: This University of Missouri corn trial revealed that Vitazyme reduced the level of most free amino acids in leaf tissue when sampled at silking, but fairly large increases in isoleucine, ornithine, and arginine offset much of the reduction. Even so, the theory of trophobiosis -- where high levels of free amino acids in tissues encourage the growth of pathogenic fungi, bacteria, nematodes, and other pests -- is somewhat substantiated by these results. Vitazyme stimulates the metabolic activity of various cycles in leaves, so proteosynthesis is encouraged to reduce the free amino acid backlog in tissues, thus reducing pest feeding potential.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Yield, Quality, and Hail Damage - University of Missouri

Researcher: Manjula Nathan, Ph. D., and Timothy Reinbott, research station supervisor

Location: University of Missouri Division of Plant Sciences, Columbia, Missouri, research farm

Variety: unknown hybrid

Planting date: May 3, 2011

Row spacing: 30 inches

Seeding rate: 32,000 seeds/acre

Soil type: unknown

Experimental design: A plot area of 12 rows – six rows for a control and six rows for the Vitazyme treatment, or 0.33 acre for each area – was selected; each row was 200 feet long. Corn was grown to determine the effects of Vitazyme on the yield and growth characteristics of corn.

1. Control

2. Vitazyme


Fertilization: station standard for corn

Vitazyme application: The corn seeds of Treatment 2 were seed treated with straight Vitazyme to give a coating of 13 oz on 32,000 seeds, or an equivalent of 13 oz/acre. The control received water only on the seeds. On June 28 (V10 stage), a soil/foliar application of 13 oz/acre was made to Treatment 2.

Hail damage event: On July 2, a severe hail storm (stones of golf ball size) struck the plot area, resulting in considerable damage to the crop, with many partially or completely broken stalks. Since there appeared to be a difference in stalk damage between the two treatments, a count was made of broken stalks.

Treatment	Broken stalks stalks/acre	Difference stalks/acre
Control	11,671	—
Vitazyme	9,365	(-) 2,306 (-20%)

**Decrease in broken stalks
with Vitazyme: -20%**



It is clear that the Vitazyme treated corn had substantially stronger stalks due to greater deposition of lignin, cellulose, and hemicellulose in the stalk structures.

Plant analysis results: On July 14, at silking, some of the hail-damaged ear-leaf tissue was collected and analyzed at the University of Missouri Soil and Crop Testing Laboratory. Results are shown below. d.m. = dry matter.

Treatment	Nitrogen % d.m.	N change % d.m.	Phosphorus % d.m.	P change % d.m.	Potassium % d.m.	K change % d.m.	Crude protein %	Protein change %
Control	2.126	—	0.290	—	1.323	—	13.3	—
Vitazyme	2.188	0.062 (+3%)	0.338	0.048 (+17%)	1.550	0.227 (+17%)	13.7	0.4 (+3%)

Increase in leaf N with Vitazyme: 3%
Increase in leaf P with Vitazyme: 17%
Increase in leaf K with Vitazyme: 17%
Increase in leaf crude protein with Vitazyme: 3% (0.4%-point)

All nutrients and protein were increased in leaf tissue harvested at silking.

Grain analysis results: At harvest, grain samples of the two treatments were analyzed at the University of Missouri Soil and Crop Testing Laboratory, with the following results.

Treatment	N	P	K	Ca	Mg	Zn	Fe	Mn	Cu	B	Mo	S	N:S
	%	%	%	%	%	ppm	ppm	ppm	ppm	ppm	ppm	%	
Control	1.20	0.35	0.36	0.01	0.06	16.8	13.2	3.9	2.6	28.2	1.0	0.007	171
Vitazyme	1.20	0.55	0.40	0.01	0.09	20.4	14.6	5.1	2.9	27.5	0.3	0.005	240

Increases with Vitazyme


P 0.20 %-point **Fe 1.4 ppm**
K 0.04 %-point **Mn 1.2 ppm**
Mg 0.03 %-point **Cu 0.3 ppm**
Zn 3.6 ppm **N:S 171 to 240**

In most cases, nutrient uptake into the grain was increased with Vitazyme application.

Yield results: Due to the July 3 hail storm, yields were greatly reduced, and are given below.

Treatment	Grain yield	Yield change
	bu/acre	bu/acre
Control	48	—
Vitazyme	56	8 (+17%)

Yield increase with Vitazyme: 17%



Conclusions: In this side-by-side experimental farm demonstration at Columbia, Missouri, a seed plus foliar treatment of Vitazyme resulted in a substantial 8 bu/acre (17%) yield increase despite a severe hailstorm on July 3. Of considerable interest is the fact that the hail storm broke off 20% fewer stalks that had been treated with this product due to stronger connective tissues – cellulose, hemicellulose, and lignin – in the treated plants. Tissue levels of N, P, K, and protein were all increased, especially P, and K, and grain elemental levels at harvest was in most cases increased, especially for P. These results reveal the ability of Vitazyme’s active agents to activate rhizosphere activity and the availability of nutrients, especially P, presumably because of a more active mycorrhizae population. An analysis of free amino acids in the plant tissues is discussed in a separate report.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Researcher: Eddie Pearson

Variety: Pioneer 2058

Seeding rate: 26,500 seeds/acre

Experimental design: Two adjacent corn fields, both treated equally and of the same soil type, were selected to compare a single Vitazyme treatment with an untreated area. A 65-acre field served as the untreated control, and an adjacent 10-acre area was treated with Vitazyme.

Location: Hooser Farm, Frost, Texas

Planting date: February 27, 2012

Row spacing: 30 inches

1. Control

2. Vitazyme

Fertilization: before planting, 150 lb/acre anhydrous ammonia (82-0-0 %N-P₂O₅K₂O), 150 lb/acre 18-46-0, and 150 lb/acre 32-0-0; during growth, 110 lb/acre 34-0-0 sidedressed

Vitazyme application: 13 oz/acre sprayed over the leaves and soil sometime between February 27 and May 10, 2012

Harvest date: July 19, 2012

Yield results: At harvest, three 30-foot typical sections of row for both treatments were picked and bagged, to make a composite 90-foot (225 ft.²) row sample for each.

Treatment	Ear number	Ear weight	Estimated grain ¹	Yield ²	Yield change
		lb	lb	bu/acre	bu/acre
Control	110	58.8	47.0	162.5	—
Vitazyme	123 (+12%)	65.8 (+12%)	52.6	181.8	19.3 (+12%)

¹A shelling percentage of 80% grain on the ear is estimated. Moisture content is estimated at 14%.

²Area harvested for each treatment = 0.0051653 acre. 1 bushel = 56 lb.

**Ear number increase
with Vitazyme....12%**

**Ear weight increase
with Vitazyme....12%**

**Yield increase with
Vitazyme....12%**

Income results: Based on corn valued at \$8.30/bu, 13 oz/acre of Vitazyme increased the value of corn produced by \$160.19/acre.

Aflatoxin levels: Samples of corn grain from each treatment were sent to A and L Laboratories in Lubbock, Texas. Both samples tested negative (<2 ppb) for aflatoxins.

Conclusions: By applying 13 oz/acre one time to this corn field in southern Texas, the yield was increased by 19.3 bu/acre, giving an increased income of about \$160/acre. This shows the efficacy of even a single application of Vitazyme for corn, although the total program recommended a 13 oz/acre application on the seeds at planting, which likely would have increased the yield even more.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Researcher: V. Plotnikov

Research organization: National Academy of Agricultural Sciences

Location: Vinnytsia, Ukraine

Variety: DKS 2960 (FAO 250)

Previous crop: corn

Seedbed preparation: plowing, harrowing, and cultivation

Soil type: gray podzolic (2.2% organic matter, 8.4 mg/100 g of soil hydrolyzed N, 15.8 mg/100 g of soil P, 12.4 mg/100 g of soil exchangeable K, pH = 5.5)

Planting date: May 14, 2012

Seeding rate: 80,000 seeds/ha

Experimental design: A small plot corn study was laid out on land previously grown to corn, using 1.1 ha plots and four replications. The purpose of the study was to determine the effect of Vitazyme biostimulant on corn growth and yield, with one treatment using soil on which a Vitazyme treated corn crop had been grown in 2011.

1. Control

2. Vitazyme

3. Vitazyme in 2011

Fertilization: 100 kg/ha N, 60 kg/ha P₂O₅, and 60 kg/ha K₂O applied before cultivation


Vitazyme application: 1 liter/ha sprayed on the leaves and soil at five to six mature leaves

Yield and income results:

Treatment	Grain yield tons/ha	Yield change tons/ha	Income increase hrn/ha
Control	9.75	—	—
Vitazyme	10.70	0.95 (+10%)	1,700
Vita in 2011	10.22	0.47 (+5%)	880

**Yield increase with Vitazyme,
1 liter/ha: 10%**

**Yield increase with Vitazyme,
applied in 2011: 5%**




Vitazyme produced an excellent 10% yield increase with 1 liter/ha applied, and there was a significant 5% increase from residual effects from 2011 applications

Leaf area results:

Treatment	Leaf area	Leaf area change
	m ² /m ² of soil	m ² /m ² of soil
Control	2.8	—
Vitazyme	3.6	0.8 (+29%)
Vita in 2011	3.1	0.3 (+11%)

Increase in leaf area with Vitazyme, 1 liter/ha: 29%

Increase in leaf area with Vitazyme, applied in 2011: 11%



A great response in leaf area was noted with Vitazyme application, 29% more leaf area with the 1 liter/ha application, and 11% more with the 2011 treatment. The carryover effect was very apparent.

Conclusions: A replicated Ukrainian corn study, using Vitazyme at the 5 to 6 leaf stage, provided a yield increase of 10%, and a consequential income increase of 1,700 hrn/ha. Corn treated on the same plots in 2011 caused a respectable 5% yield increase. Corn leaf area was also increased by Vitazyme (by 29%) for the 1 liter/ha application, and by 11% with the 2011 treatments. These results show the strong carryover effect of the product, and the highly positive and profitable yield results, making it an excellent choice for farmers in Ukraine.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Researcher: Bert Schou, Ph.D.

Research organization: Agricultural Custom Research and Education Services (ACRES)

Location: Cedar Falls, Iowa Variety: Pioneer PO 448 (non-GMO) Previous crop: soybeans

Soil type: Aredale Loam (36% sand, 42% silt, 22% clay), 3.6% organic matter, pH = 6.1, C.E.C. = 15.2 meq/100 g, fertility level = excellent, soil drainage = excellent Planting depth: 1.5 inches

Row spacing: 30 inches Planting rate: 38,000 seeds/acre Seedbed at planting: fine

Planting date: May 9, 2012 Plot size: 15 x 40 feet (600 ft²) Tillage: conventional

Irrigation: 2 inches total in late July and early August

Experimental design: A small plot, replicated corn trial (four replicates) was established in eastern Iowa to evaluate the ability of two Vitazyme variations, plus an amino acid formulation, alone or in combination, to influence corn yield and quality.

Treatment	Vitazyme 1*	Vitazyme 2*	Amino acids*
1	0	0	0
2	13 oz/acre (2x)	0	0
3	0	13 oz/acre (2x)	0
4	0	0	2 oz/acre (2x)
5	13 oz/acre (2x)	0	2 oz/acre (2x)

*All applications were on the seeds at planting, and on the leaves and soil later.

Fertilization: Fall of 2011, 60 lb/acre of 18-46-0% N-P₂O₅-K₂O; June 5, 2012, 110 lb/acre of N as urea ammonium nitrate, with a sprayer and drop nozzles

Vitazyme application: For Treatments 2, 3, and 5, 13 oz/acre on the seeds at planting (May 9), and again at 13 oz/acre on the leaves and soil at the V6 stage (June 16). Treatments 2 and 5 received Vitazyme 1, and Treatment 3 received a slight modification called Vitazyme 2.

Amino acid application: A proprietary amino acid blend was applied to Treatments 4 and 5, to the seeds at 2 oz/acre on May 9, and to the leaves and soil at the V6 stage at 2 oz/acre on June 16. For Treatment 5, the amino acids were mixed with the Vitazyme before application.

Sprayer settings: seed treatment, 10 gallons/acre of 115 ml of Vitazyme in 3 gallons of water, or 18 ml of amino acids in 3 gallons of water; foliar and soil treatment, 15 gallons/acre of 77 ml of Vitazyme in 3 gallons of water, or 12 ml of amino acids in 3 gallons of water


Weed control: preplant: Harness Xtra at 1.2 quarts/acre; additional post-emergence applications

Weather during the growing season: The season was hot and dry, with the July average high temperature being 92.3° F, and the August average high being 85.5° F. Rainfall for April through October 12 was 12 inches, whereas the normal is 26 inches.

Harvest date: October 20, 2012. A Massey-Ferguson 9 plot combine harvested the middle two rows of each plot, and the corn was weighed on an electronic scale.


Plant population results: No significant differences were detected.

Corn test weight: Although differences among treatments were not significant, all four Vitazyme and amino acid treatments exceeded the control.




Corn protein and fat: Composite corn samples from the four replicates of each treatment were sent to Midwest Laboratories, Inc., Omaha, Nebraska, to evaluate the levels of protein and fat in the samples. Fat of the treated corn always exceeded the untreated control, and in all but one case the protein was also evaluated by the treatments.

Protein (dry basis)



Fat (dry basis)



Increase grain protein with Vitazyme and amino acids

Vitazyme 2	+2%
Amino acids	+1%
Vitazyme 1 + Amino acids	+3%

Increase grain fat with Vitazyme and amino acids

Vitazyme 1	+3%
Vitazyme 2	+7%
Amino acids	+20%
Vitazyme 1 + Amino acids	+3%


Changes in grain composition are usually quit resistant to treatment, but there were small increases in grain protein with Vitazyme and amino acids. Fat composition was improved by all four treatments, and dramatically by the amino acid and Vitazyme 1 plus amino acid treatments (19 to 20%).

Yield results: The two inner rows of each plot were harvested.

Treatment	Yield ¹ bu/acre	Yield change bu/acre
1. Control	199.10 b	—
2. Vitazyme 1	223.44 a	24.34 (+12%)
3. Vitazyme 2	213.76 ab	14.66 (+7%)
4. Amino acids	215.18 ab	16.08 (+8%)
5. Vita 1 + A.A.	208.28 ab	9.18 (+5%)
LSD _{0.05}	15.21 bu/acre	
Standard deviation	11.35 bu/acre	
Replicate F	10.947	
Treatment F	3.150	
C.V.	5.35%	

¹Means followed by the same letter are not significantly different at P = 0.05, according to the Student-Newman-Keuls Test.

Yield increase with Vitazyme and amino acids	
Vitazyme 1	+12%
Vitazyme 2	+7%
Amino acids	+8%
Vitazyme + Amino acids	+5%



Notice that Vitazyme 1 increased the corn yield by 12% (24.34 bu/acre), while the other treatments boosted the yield by from 5 to 8% above the control. While these lower increases were not significant at P = 0.05, yet the 7 and 8% increases were very close to being significant.

Conclusions: A replicated corn study in east-central Iowa, during a very hot and dry year, revealed that two variations of Vitazyme both increased corn yield, one by 12% (24.34 bu/acre), and another by 7% (14.66 bu/acre). An amino acid product increased yield by 8% (16.08 bu/acre). With a corn price of \$7.50/bu during the fall of 2012, these yield increases translate to \$182.55, \$109.95, and \$120.60/acre. The combined Vitazyme plus amino acid treatment increased yield a very respectable 5% (9.18 bu/acre), but did not display a synergism. Grain quality was also improved by both Vitazyme and amino acids. Per bushel weight was increased by up to 0.51 lb/bu (Vitazyme 2), and grain protein was improved by 1 to 3%. Grain fat was especially enhanced by both products, increasing by 3 to 7% with Vitazyme alone, and by 20% with amino acids alone, and by 19% with the two products combined. These results show the great value of utilizing Vitazyme, and to a lesser extent the amino acids, to improve corn grain yield and quality in the central Corn Belt. There appears to be no synergism between Vitazyme and the amino acid formulation used in this study.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

A Greenhouse Study

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Planting date: January 19, 2012

Pot size: 1 gallon

Planting rate: 8 seeds/pot, thinned to 3 plants/pot

Soil type: silt loam

Experimental design: A greenhouse pot trial using 12 replicates was set up to evaluate the effect of Vitazyme, applied once at a standard field rate, on corn growth.

1. Control

2. Vitazyme

Fertilization: none

Vitazyme application: 100 ml of a 0.1% solution for each pot, immediately after planting

Harvest date: March 5, 2012, 46 days after planting


Plant height results: All plants were measured to the nearest cm, and the three plants for each pot were averaged.

Treatment	Plant height cm	Height change cm
Control	95.7 a	—
Vitazyme	99.3 a	3.6 (+4%)
Block P	0.382	
Treatment P	0.153	
Model P	0.336	
CV _{0.10}	5.85%	
LSD _{0.10}	4.2 cm	

Vitazyme increased corn height by 4% at 46 days after planting, but this increase was not significant at P = 0.10.

Dry weight results: The plants were placed in a drying oven for 24 hours at 115°F, and then weighed to the nearest 0.01 gram.

Treatment	Dry Weight	Weight change
	grams	grams
Control	15.27	—
Vitazyme seed trt.	18.33	3.06 (+20%)
Block P	0.7148	
Treatment P	0.0029**	
Model P	0.1589	
CV _{0.10}	11.74%	
LSD _{0.10}	1.45 grams	



Increase in dry weight with Vitazyme: 20%

Vitazyme dramatically and significantly increased the dry matter accumulation of the corn with only one application, by 20%.

Conclusions: In this greenhouse trial, Vitazyme greatly increased the fixation of dry matter in the plants with 100 ml/pot of a 0.1% solution applied at planting. The increase was highly significant, and 20% greater than the control. Plant height was increased by 4% with Vitazyme, but this increase was not significant. These results verify the utility of this product, with its natural vitamins and growth regulators, to improve plant growth and yield.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

Seed Treatment Results in the Greenhouse

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Planting date: March 14, 2012

Pot size: 1 gallon

Soil type: silt loam

Seeding rate: 8 seeds/pot, thinned to 3 plants/pot

Experimental design: A four replicated greenhouse trial, using corn seeds pre-treated with Vitazyme on January 17, 2012, was planted on March 14, 2012, 57 days after treatment, to determine residual effects of the seed treatment on plant growth. A randomized complete block design was used.

1. Control

2. Vitazyme seed treatment


Fertilization: none

Vitazyme treatment: On January 17, corn seeds were soaked in a 10% Vitazyme solution for 6 minutes, then dried on paper towels with a fan blowing across them for rapid drying. The seeds were stored in closed jars at room temperature along with untreated seeds, which served as the control.

Harvest date: April 5, 2012, 21 days after planting

Dry weight results: Roots of the corn plants were washed free of soil, and the plants were placed in a drying oven at 115 °F for 24 hours. Weights of the dried plants for each pot were recorded to the nearest 0.01 gram.

Treatment	Dry Weight	Weight change
	g	g
Control	3.12	—
Vitazyme seed trt.	3.55	0.43 (+14%)
Block P	0.339*	
Treatment P	0.0978*	
Model P	0.0403*	
CV _{0.10}	7.581%	
LSD _{0.10}	0.42 gram	



**Increase in dry weight with
Vitazyme: 14%**

A seed treatment on these corn seeds 57 days before planting caused a significance 14% increase in dry weight at harvest.

Conclusions: A greenhouse pot study, using corn seeds treated with a 10% Vitazyme solution and dried 57 days earlier, produced a 14% increase in dry weight after 21 days of growth. This significant increase at $P = 0.10$ shows the effectiveness of a seed treatment to improve corn growth, and the long-term efficacy of the practice.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Corn

A Greenhouse Trial -- Synergism With Amino Acids

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Planting date: January 19, 2012

Pot size: 1 gallon

Planting rate: 8 seeds/pot, thinned to 3 plants/pot

Soil type: silt loam

Experimental design: A replicated greenhouse pot study (four reps) use set up to evaluate the effects of Vitazyme and certain amino acids — alone and together — on corn growth.

1. Control

2. Vitazyme

3. Amino acids

4. Vitazyme + Amino acids

Fertilization: none

Vitazyme application: Pots of Treatments 2 and 4 received 100 ml/pot of a 0.1% Vitazyme solution just after seeding.


Amino acid application: A special liquid proprietary amino acid blend was applied at 100 ml of a 0.04% solution to Treatments 3 and 4. For Treatment 4, the product was mixed with Vitazyme in 100 ml of water.

Harvest date: March 1, 2012, 41 days after planting

Height results: The three plants from each pot were measured to the nearest cm, and averaged. A statistical analysis was then performed on these averages.

Plant Height

Treatment	Plant height cm	Height change cm
Control	82.6 b	—
Vitazyme	91.9 a	9.3 (+11%)
Amino acids	92.7 a	10.1 (+12%)
Vita + A.A.	88.4 a	5.8 (+7%)
Block F	0.0096*	
Main effects F	0.0262*	
Model F	0.0088*	
LSD _{0.10}	5.3 cm	
CV _{0.10}	4.63%	




Increase in height with Vitazyme: 11%
Increase in height with Amino Acids: 12%

All three treatments significantly increased plant height at P = 0.10.

Dry weight results: The harvested plants were dried in a drying oven for 24 hours at 115°F, before weighing to the nearest 0.01 gram.

Plant Dry Weight



Treatment	Dry weight	Weight change
	g	g
Control	7.39 b	—
Vitazyme	9.32 a	1.93 (+26%)
Amino acids	9.50 a	2.11 (+29%)
Vita + A.A.	9.80 a	2.41 (+33%)
Block F	0.184	
Main effects F	0.068*	
Model F	0.088*	
LSD _{0.10}	1.54 gram	
CV _{0.10}	13.2%	

Increase in dry weight

Vitazyme	26%
Amino acids	29%
Vitazyme + Amino Acids	33%

All three treatments increased plant dry weight at P = 0.10, especially the combined Vitazyme + Amino Acid treatment.

Conclusions: In this greenhouse study with corn, Vitazyme, Amino Acids, and the combined products all significantly increased both plant height and dry weight. Dry weight was especially

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
 (903) 845-2163 FAX: (903) 845-2262

2012 Crop Results

Vitazyme on Baby Corn

Researcher: Richard Stonewigg

Research organization: Lachlan Kenya Limited.

Location: Mboga Tuu, Isinya, Kenya

Soil type: unknown

Variety: “Baby” corn

Planting date: November, 2011

Irrigation: drip

Experimental design: A field of baby corn was divided into five treatments of 1,800 m² each to evaluate the effect of Vitazyme (two formulations) and Rizobacter, alone and in combination, on the health, growth parameters, and the yield of the crop.

- 1. Control
- 2. Vitazyme, regular
- 3. Rizobacter
- 4. Vitazyme, regular + Rizobacter
- 5. Vitazyme, from concentrate + Rizobacter

Fertilization: unknown


Vitazyme treatment: The Vitazyme from concentrate was diluted with water (1 part concentrate to 9 parts water). Treatments for either product alone, or combined with Rizobacter, were as follows:

Time	Rate
Seeding	5 ml/kg of seed
Drench, pre-plant	200 ml/20 liters of water
Two weeks post-plant	foliar 50 ml/20 liters of water
Before tasseling	foliar 50 ml/20 liters of water
At tasseling	foliar 50 ml/20 liters of water
At ear filling	foliar 50 ml/20 liters of water

Rizobacter application: The Rizobacter products are bacterial preparations from Argentina.

Yield results: The crop was harvested in January of 2012.

Treatment	Yield tons	Yield increase tons
1. Control	0.490	—
2. Vitazyme, regular	0.935	0.445 (+91%)
3. Rizobacter	0.890	0.400 (+82%)
4. Vita (regular) + Rizobacter	1.024	0.534 (+109%)
5. Vita (from conc.) + Rizobacter	1.113	0.623 (+127%)



Yield increase with Vitazyme

- Vitazyme regular +91%**
- Vitazyme regular + Rizobacter +109%**
- Vitazyme concentrate + Rizobacter +127%**

Conclusions: This baby corn study in Kenya proved that both Vitazyme and Rizobacter inoculant gave excellent yield increases, especially when the two products were combined. Regular Vitazyme gave a 91% yield increase, while Rizobacter alone provided an 82% increase. When regular Vitazyme was combined with Rizobacter, a 109% yield increase resulted, showing the synergism between the bacterial product and Vitazyme. Both the regular and the diluted concentrate (10:1) provided excellent yield increases (109% and

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher: Unknown

Research coordinator: I.V. Braginets

Research organization: Alfa-Agro, Ukraine

Variety: unknown

Experimental design: A field was divided into a Vitazyme treated and an untreated portion to evaluate the effect of this product on crop yield.

1. Control

2. Vitazyme

Fertilization: farm practice

Vitazyme application: 1 liter/ha sprayed on the leaves and soil at the 10 to 12-leaf stage

Yield results: No yield results are available, but the increase in yield is given.

**Increase in corn yield with Vitazyme:
1.72 tons/ha (27.4 bu/acre)**

Conclusion: This yield increase was an excellent result of Vitazyme application in this Ukraine study.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher/Farmer: Nong Van Duc
Nong Province, Viet Nam

Location: Village 7, Eapo Hamlet, Cu Jut District, Dak
Variety: NK 7328

Planting season: Summer – Fall, 2011

Experimental design: A corn field was divided into a Vitazyme treated area of 1.0 ha, and an untreated control area of 0.5 ha, to determine the effect of this product on the yield and profitability of the crop.

1. Control

2. Vitazyme

Vitazyme application: (1) a 5% Vitazyme seed spray on the seeds before planting; (2) 1 liter/ha sprayed on the leaves and soil 35 days after planting


Growth results: With Vitazyme the following effects were noted:

- Taller plants
- Stronger root systems
- Darker green leaves
- Excellent disease resistance from steaked leaf and stem rot
- During a drought period, plants showed no leaf rolling
- Seed color was brighter

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	8.6	—
Vitazyme	9.7	1.1 (+13%)

**Increase in corn yield with
Vitazyme: 13%**



Income results:

Treatment	Vitazyme ¹ VND/ha	Total costs VND/ha	Total income ² VND/ha	Net income VND/ha	Extra profit VND/ha
Control	0	11,250,000	38,700,000	26,700,000	—
Vitazyme	500,000	11,500,000	43,650,000	31,150,000	4,450,000

¹VND = Vietnamese dollar; 1 USD = 20,000 VND.

²Corn price = 4,500 VND/kg.

**Increase in income with Vitazyme:
4,450,000 VND, or \$222.50/ha**

Conclusion: This Vietnamese corn trial revealed that Vitazyme, applied on the seeds as well as 35 days after planting, increased yield by 13%, while boosting profit by \$222.50/ha. These results prove the great utility of the Vitazyme program to grow corn in Viet Nam.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher/Farmer: Dinh Thi Ngan
Province, Viet Nam

Location: Village 2, Dak Will, Cu Jut District, Dak Nong
Variety: NK 72

Planting season: Summer – Fall, 2011

Experimental design: A corn field was divided into a Vitazyme treated area of 0.9 ha, and an untreated control area of 0.5 ha, to determine the effect of this product on the yield and profitability of the crop.

1. Control

2. Vitazyme

Vitazyme application: a 5% Vitazyme seed spray on the seeds before planting


Growth results: With Vitazyme the following effects were noted:

- Taller plants
- Stronger root systems
- Darker green leaves
- Excellent disease resistance from steaked leaf and stem rot
- During a drought period, plants showed no leaf rolling
- Seed color was brighter

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	8.0	—
Vitazyme	9.0	1.0 (+13%)

**Increase in corn yield with
Vitazyme: 6%**



Income results:

Treatment	Vitazyme ¹ VND/ha	Total costs VND/ha	Total income ² VND/ha	Net income VND/ha	Extra profit VND/ha
Control	0	12,000,000	36,000,000	24,750,000	—
Vitazyme	250,000	12,250,000	38,250,000	27,000,000	2,250,000

¹VND = Vietnamese dollar; 1 USD = 20,000 VND.

²Corn price = 4,500 VND/kg.

**Increase in income with Vitazyme:
2,250,000 VND, or \$112.75/ha**

Conclusion: This Vietnamese corn trial revealed that Vitazyme, applied as a 5% seed treatment at planting, increased yield by 6%, while boosting profit by \$112.75/ha. Had an additional foliar treatment been applied, yields and profits would likely have improved even more.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher/Farmer: Tran Van Nhung
Nong Province, Viet Nam

Location: Village Tan Thanh, Eapo, Cu Jut District, Dak
Variety: SSC 557

Planting season: Summer – Fall, 2011

Experimental design: A corn field was divided into a Vitazyme treated area of 0.5 ha, and an untreated control area of 0.2 ha, to determine the effect of this product on the yield and profitability of the crop.

1. Control

2. Vitazyme

Vitazyme application: 1.5 liters/ha sprayed on the leaves and soil 30 days after planting


Growth results: With Vitazyme the following effects were noted:

- Taller plants
- Stronger root systems
- Darker green leaves
- Excellent disease resistance from steaked leaf and stem rot
- During a drought period, plants showed no leaf rolling
- Seed color was brighter

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	7.4	—
Vitazyme	8.0	0.6 (+8%)

**Increase in corn yield with
Vitazyme: 8%**



Income results:

Treatment	Vitazyme ¹ VND/ha	Total costs VND/ha	Total income ² VND/ha	Net income VND/ha	Extra profit VND/ha
Control	0	11,125,000	31,080,000	19,955,000	—
Vitazyme	575,000	11,700,000	33,600,000	21,900,000	1,945,000

¹VND = Vietnamese dollar; 1 USD = 20,000 VND.

²Corn price = 4,200 VND/kg.

**Increase in income with Vitazyme:
1,945,000 VND, or \$97.25/ha**

Conclusion: This Vietnamese corn trial revealed that Vitazyme, applied once at 1.5 liters/ha at 30 days after planting, increased yield by 8%, while boosting profit by \$97.25/ha. Had a seed treatment been applied, yields and profits would likely have improved even more.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

Average Values for 2009 to 2011 in Ukraine

Vitazyme on Corn

Researcher: V.V. Plotnikov


Location: National Academy of Agrarian Sciences, Vinnytsia State Agricultural Research Station, Vitnnytsia, Ukraine (Central Forest and Steppe Region)

Demonstration plot values averaged over three years, 2009 to 2011:

Treatment	Yield tons/ha	Yield change tons/ha
1. Control	8.85	—
2. Vitazyme on leaves ¹	9.88	1.03 (+12%)

¹1 liter/ton on the leaves + soil at the 7 to 8-leaf stage.

**Three-Year Average Increase
With Vitazyme: +12%**



Conclusion: Over three years of demonstrations, Vitazyme is shown to be an excellent adjunct to corn production in Ukraine.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher: Michael Rethwisch
Extension, David City, Nebraska

Variety: Kruger K-4510

Experimental design: A corn field was divided into replicated strips with four different products, with Vitazyme applied to the side of the seeds, to evaluate the yield and test weight responses. All treatments received starter fertilizer under conventional tillage.

Research organization: University of Nebraska-Lincoln

Location: Aurora, Nebraska

Soil type: unknown


Treatment	Starter fertilizer	Product rate amount/acre
1. Control	X	—
2. CALFA	X	10 oz
3. Foliar Blend	X	16 oz
4. Torque	X	8 oz
5. Vitazyme	X	13 oz

Fertilization: Besides typical N-P₂O₅-K₂O fertilizer, all products and the control received 5 gallons/acre of NaChurs 6-24-6 Starter fertilizer applied to the side of the seeds with a “splitter.”

Vitazyme and other product applications: At planting, all four products were applied at the rates shown in the above table, using the “splitter.”

Weather for the growing season: Rains were ample to excessive, and temperatures were somewhat below normal.


Test weight results: Grain volume weights were determined corrected to 15.5% moisture.



While none of the test weight values were significantly different, Vitazyme had the highest test weight, which was 0.3 lb/bu greater than the closest other product value.

**Increase in test weight with
Vitazyme: 0.8 lb/bu**

Yield results:



Yield values were not significantly different among the five treatments, but Vitazyme produced the greatest yield, of 1.5 bu/acre above the control.

Increase in yield with Vitazyme: 1.5 bu/acre

Conclusion: In this eastern Nebraska corn study using four products applied at planting, using a “splitter”, Vitazyme produced the highest yield and the heaviest test weight, although the values did not differ significantly. The reason for a less than significant response with Vitazyme may have been because the product was applied to the side of the seeds, not directly on them.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn
A Nematode Evaluation

Researcher: Michael Rethwisch
David City, Nebraska

Research organization: University of Nebraska - Lincoln Extension,
Location: Silver Creek, Nebraska

Variety: Mycogen 2T832

Soil type: Darr sandy loam

Experimental design: A replicated corn trial having Vitazyme applied twice, plus the normal farming program, was evaluated for nematodes and statistically analyzed.

1. Control

2. Vitazyme

Fertilization: normal farming program

Vitazyme application: 13 oz/acre (1 liter/ha) in the seed row with starter fertilizer at planting

Growing season weather: excessive rain and normal temperatures

Nematode results: After harvest in the fall of 2010, soil samples were taken in the vicinity of the roots of each plot and analyzed at the nematode testing facilities of the University of Nebraska at Lincoln.

Treatment	Nematode species ¹		
	Stunt ²	Lesion ³	Dagger ⁴
Control	119.5	119	711.8
Vitazyme	133.0	93	352.3 (-149%)
p-value (0.05)	0.61	0.36	0.0016**


¹All data from each plot was subjected to statistical analyses to arrive at probabilities, as shown here.

²*Tylenchorhynchus* spp. They can possibly cause stunting of corn, and feed externally on the roots.

³*Pratylenchus* spp. These can be important root pathogens causing stunting of corn plants. They feed inside roots.

⁴*Xiphinema* spp. This nematode feeds externally on roots, and can survive sandy soils but is sensitive to tillage.

Dagger Nematodes



Reduction in dagger nematodes with Vitazyme: 149%

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

An Evaluation of Soil Phosphorus Levels and Two Formulations

Researcher: Bert Schou, Ph.D. **Research organization:** Agricultural Custom Research and Education Services (ACRES)
Location: Cedar Falls, Iowa **Variety:** Pioneer P0528 (non-GMO)
Soil type: Kenyon loam (34% sand, 46% silt, 20% clay, 3.6% organic matter, pH = 7.2, C.E.C. = 15 meq/100 g, fertility level = excellent, drainage = excellent) **Planting depth:** 2 inches
Row spacing: 30 inches **Planting rate:** 35,000 seeds/acre **Seedbed at planting:** fine
Planting date: May 16, 2011 **Tillage:** conventional **Plot size:** 15 x 40 feet (600 ft.²)
Experimental design: A small plot, six replicate study with corn, using three phosphorus levels in a phosphorus-deficient soil, received nine treatments, with two Vitazyme formulations. The trial was conducted to determine effects on corn yield and grain quality.

Treatment	Phosphorous lb/acre	Vitazyme A ¹ oz/acre	Vitazyme B ¹ oz/acre
1	0	0	0
2	0	13 (2X)	0
3	0	0	13 (2X)
4	65	0	0
5	65	13 (2X)	0
6	65	0	13 (2X)
7	130	0	0
8	130	13 (2X)	0
9	130	0	13 (2X)

¹2X = two applications.

Fertilization: All plots received 100 lb/acre of nitrogen and 75 lb/acre of K₂O preplant, plus the P₂O₅ applications of 0, 65, or 130 lb/acre as shown in the table above

Weed control: Harness at 1.2 qt/acre preemergence, giving excellent weed control

Vitazyme application: (1) 13 oz/acre (1 liter/ha) in the seed row at planting on May 16, 2011; (2) 13 oz/acre (1 liter/ha) on the leaves and soil at V7 on June 30, 2011. Vitazyme A and B are different formulations of Vitazyme.

Weather during the growing season: The season was favorable to the corn growth in terms of temperature and rainfall, although early July was quite dry.


Harvest date: October 15, 2011. A Massey-Ferguson 9 plot combine harvested the middle two rows of each plot, and the corn was weighed using an electronic scale.

Plant Population: There were no significant differences among the nine treatments, so these data are not


included here.

Tissue test results: On July 12, 2011, tissue analyses were conducted on composite samples for each treatment at Perry Agricultural Laboratory, Bowling Green, Missouri. The results for the first three treatments (no added phosphorus) are presented here, to reveal effects of Vitazyme A and Vitazyme B on tissue element levels.


Nitrogen




Phosphorus




Potassium




Calcium




Magnesium




Sulfur




Boron




Iron




Manganese



Copper



Zinc



Notice that in most cases Vitazyme A and Vitazyme B increased the elemental composition of the corn tissue. For potassium, sulfur, boron, manganese, and copper there were either equal contents or a lower content of the element for one product or the other.

Test weight results: There were some differences in grain test weight.


Treatment	Test weight ¹	Test weight change ²
	lb/bu	lb/bu
1 (O P)	55.17 b	—
2 (O P, Vita A)	55.82 ab	0.65 (+1%)
3 (O P, Vita B)	55.57 ab	0.40 (+1%)
4 (Low P)	56.39 a	—
5 (Low P, Vita A)	55.94 ab	(-) 0.45 (-1%)
6 (Low P, Vita B)	56.25 ab	(-) 0.14 (0%)
7 (High P)	56.04 ab	—
8 (High P, Vita A)	56.48 a	0.44 (+1%)
9 (High P, Vita B)	56.55 a	0.51 (+1%)
LSD _{0.05}	0.77 lb/bu	
Standard deviation	0.66 lb/bu	
Replicate F	4.21	
Treatment F	2.86	
CV	1.18%	

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.

²Comparisons are made with the control at the same phosphorus level.

Combined Phosphorus Test Weight for All Treatments

P level	Test weight	Test weight change
lb/acre	lb/bu	lb/bu
0	55.52	—
65	56.19	+0.67
130	56.36	+0.84




Note the trend towards higher test weight with higher P levels.

Combined Product Test Weight for All Treatments

Product	Test weight	Test weight change
	lb/bu	lb/bu
None	55.52	—
Vitazyme A	56.08	+0.21
Vitazyme B	56.12	+0.25

Both Vitazyme A and Vitazyme B show increased test weights versus the untreated controls. This result is likely due to the products' enhanced rhizosphere activation so more nutrients were available for uptake.




Yield results:

Treatment	Yield ¹ bu/acre	Yield change ² bu/acre
1 (O P)	187.1 c	—
2 (O P, Vita A)	190.3 bc	3.2 (+2%)
3 (O P, Vita B)	197.7 abc	10.6 (+6%)
4 (Low P)	196.7 abc	—
5 (Low P, Vita A)	195.8 abc	(-) 0.9 (0%)
6 (Low P, Vita B)	199.8 abc	3.1 (+2%)
7 (High P)	202.6 ab	—
8 (High P, Vita A)	203.3 ab	0.7 (0%)
9 (High P, Vita B)	209.9 a	6.6 (+3%)
LSD _{0.05}	9.7 bu/acre	
Standard deviation	8.3 bu/acre	
Replicate F	3.27	
Treatment F	4.13	
CV	4.19	

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.
²Comparisons are made with the control at the same phosphorus level.

Combined Phosphorus Yield for All Treatments

P level lb/acre	Yield lb/bu	Yield change lb/bu
0	191.7	—
65	197.4	+5.7 (+3%)
130	205.3	+13.6 (+7%)





With increasing phosphorus levels the yield of corn grain increased almost in a linear fashion, to 3 and 7% above the control at 65 and 130 lb/acre of phosphorus, respectively.

Combined Product Test Weight for All Treatments

Product	Yield lb/bu	Yield change lb/bu
None	195.5	—
Vitazyme A	196.5	+1.0 (+1%)
Vitazyme B	202.5	+7.0 (+4%)

Vitazyme A, across all phosphorus levels, slightly increased corn yields, but Vitazyme B increased yields by 4%, or 7.0 bu/acre.



There was a trend across the treatments for higher yields due to higher fertilizer phosphorus applications and Vitazyme B. The highest overall treatment was combined high phosphorus plus Vitazyme B (Treatment 9), which increased yield by 22.8 bu/acre (+12%) above the untreated, low phosphorus control (Treatment 1).

Grain protein: A composite grain sample from each of the six replicates for the nine treatments was sent to Midwest Laboratories in Omaha, Nebraska, for the analysis of protein and elements. The various elements showed no trends among the treatments, but protein displayed interesting results.

Treatment	Crude protein ¹
	%
1 (O P)	6.42
2 (O P, Vita A)	6.80
3 (O P, Vita B)	6.66
4 (Low P)	6.80
5 (Low P, Vita A)	6.90
6 (Low P, Vita B)	6.60
7 (High P)	6.22
8 (High P, Vita A)	7.21
9 (High P, Vita B)	6.28

¹Dry weight basis

Combined Phosphorus Protein for All Treatments

P level	Protein	Protein change
lb/acre	%	%
0	6.63	—
65	6.77	+0.14
130	6.57	-0.06

There was no clear relationship between fertilizer and soil phosphorus and grain protein.

Combined Product Protein for All Treatments

Product	Protein	Protein change
	%	%
None	6.48	—
Vitazyme A	6.97	+0.49
Vitazyme B	6.51	+0.02


Note that Vitazyme A improved grain protein by near 0.5%, although Vitazyme B had little effect.

Conclusion: A replicated small-plot study in east-central Iowa, using two Vitazyme versions with treatments at planting and V-7, and three phosphorous fertilizer levels, revealed several significant effects of phosphorous and Vitazyme on test weight, yield, and grain protein. The year was favorable for high yields, so yields varied from 187.1 to 209.9 bu/acre. The tissue element levels for the first three treatments, analyzed mid-season, revealed that both Vitazyme A and Vitazyme B increased element levels, with few exceptions.

Test weight showed a positive impact from fertilizer phosphorous, increasing by 0.67 lb/bu for 65 lb/acre of P₂O₅, and by 0.84 lb/bu for 130 lb/acre of P₂O₅. Both Vitazyme A and Vitazyme B gave increases in the test weight, of 0.21 and 0.25 lb/bu, respectively. It is interesting that the highest test weights were for Treatments 8 and 9 (high P₂O₅, Vitazyme A and Vitazyme B, respectively), which produced significantly heavier grain than the low-P₂O₅, no-Vitazyme control (Treatment 1). This result reveals that higher phosphorous levels plus Vitazyme increase mineral uptake the most due to rhizosphere activation – especially phosphorous-extracting mycorrhizae – and increased availability of soil nutrients.

Yield results showed significant increases in corn grain with the high P₂O₅ + Vitazyme B treatment, which exceeded the no P₂O₅ and no P₂O₅ + Vitazyme A treatments. Of particular interest is the trend for increasing grain yield with both increasing phosphorous fertility and Vitazyme A and B, especially with Vitazyme B, which boosted corn yield by 7.0 bu/acre (4%) above the combined average of the no-Vitazyme phosphorous treatments at all P₂O₅ levels. The highest yield was for Treatment 9 – high P₂O₅ + Vitazyme – which boosted grain yield by 22.8 bu/acre over the no P₂O₅ + no Vitazyme control.

Grain protein responded with no pattern to added P₂O₅ fertilizer, but Vitazyme A increased protein by 0.49%-point across all three P₂O₅ fertilizer levels. This study reveals that, of the two Vitazyme formulations, Vitazyme B is the best in terms of test weight and yield enhancement, but Vitazyme A increased grain protein the most. Both products, however, improved yield, protein, and test weight of corn.



Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher: Paul W. Syltie, Ph.D. Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas
Variety: yellow dent Pot size: 1 gallon Soil type: silt loam
Planting rate: 10 seeds/pot, thinned to 3 plants/pot Planting date: November 12, 2010
Experimental design: A replicated greenhouse pot study was conducted to evaluate the effect of two Vitazyme formulations to increase plant growth. Seven replicates were used.

1. Control

2. Vitazyme A

3. Vitazyme B


Fertilization: none

Vitazyme application: Vitazyme A: 100 ml/pot at planting of a 0.1% solution, for regular product; Vitazyme B: 100 ml/pot at planting of a 0.1% solution, diluted directly from concentrate

Plant height results: The plants were measured on December 15, 2010, and averaged for each pot.

Treatment	Plant height ¹ cm	Height change cm
Control	68.9 b	—
Vitazyme A	73.0 a	4.1 (+6%)
Vitazyme B	76.1 a	7.2 (+10%)
Treatment P	0.0023**	
Model P	0.0023***	
CV	4.45%	
LSD _{0.05}	3.6 cm	

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls-Test.




**Increase in height with
Vitazyme A: 6%**

**Increase in height with
Vitazyme B: 10%**

Dry weight results: On December 15, 2010, the roots were washed free of soil, and dried in a drying oven at 120° F for 24 hours.

Treatment	Dry Weight ¹	Weight change
	grams	grams
Control	4.23 b	—
Vitazyme A	4.89 a	0.66 (+16%)
Vitazyme B	5.04 a	0.81 (+19%)
Treatment P	0.0006***	
Model P	0.0006***	
CV	7.06%	
LSD _{0.10}	0.37 gram	

¹Means followed by the same letter are not significantly different at P=0.10 according to the Student-Newman-Keuls-Test.



**Increase in dry weight with
Vitazyme A: 16%**

**Increase in dry weight with
Vitazyme B: 19%**

Conclusion: A replicated pot trial in the greenhouse, using two variations of Vitazyme, proved that both corn height and dry weight responded significantly to both products. Height increased by 6% and 10%, respectively, for Vitazyme A and Vitazyme B, while dry weight increased by 16% and 19% for the two products. These results prove that both products are very effective for improving corn growth, especially Vitazyme B, the product made directly from concentrate.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

A Greenhouse Study

Researcher: Paul W. Syltie, Ph.D. Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas
Variety: yellow dent Pot size: 1 gallon Soil type: silt loam
Planting rate: 10 seeds/pot, thinned to 3 plants/pot Planting date: February 14, 2011
Experimental design: A replicated greenhouse pot study was conducted to evaluate the relative effectiveness of two Vitazyme formulations to increase plant growth. Eight replications were used.

1. Control

2. Vitazyme A

3. Vitazyme B


Fertilization: none

Vitazyme application: Vitazyme A: 100 ml/pot at planting of a 0.1% solution, from regular product; Vitazyme B: 100 ml/pot at planting of a 0.1% solution, diluted directly from concentrate

Plant height results: On March 22 at harvest, 36 days after planting, the plants were measured for height, and averaged for the three plants in each pot.

Treatment	Plant height ¹ cm	Height change cm
Control	73.6 a	—
Vitazyme A	75.6 a	2.0 (+3%)
Vitazyme B	74.3 a	0.7 (+1%)
Block P	0.605	
Treatment P	0.463	
Model P	0.624	
CV _{0.10}	4.07%	
LSD _{0.10}	2.9 cm	

¹Means followed by the same letter are not significantly different at P=0.10 according to the Student-Newman-Keuls-Test.




Increase in plant height with Vitazyme

Vitazyme A 3%
Vitazyme B 1%

Dry weight results: On March 22 the roots were washed of all soil, and then dried in a drying oven at 120° F for 24 hours. Weights were recorded to the nearest 0.01 gram for the combined three plants of each pot.

Treatment	Dry weight ¹ grams	Weight change grams
Control	9.28 b	—
Vitazyme A	9.95 a	0.67 (+7%)
Vitazyme B	9.75 a	0.47 (+5%)
Block P	0.018*	
Treatment P	0.011*	
Model P	0.008**	
CV _{0.10}	3.64%	
LSD _{0.10}	0.34 gram	

¹Means followed by the same letter are not significantly different at P=0.10 according to the Student-Newman-Keuls-Test.



Increase in dry weight with Vitazyme

Vitazyme A 7%
Vitazyme B 5%

Conclusion: This replicated greenhouse study with corn proved that both Vitazyme A (regular product) and Vitazyme B (direct dilution from concentrate) increased dry matter production significantly above the untreated control.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
 (903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn
A Concentration Series Study


Researcher: Paul W. Syltie, Ph.D. Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas
Variety: yellow dent Pot size: 1 gallon Soil type: silt loam
Planting rate: 10 seeds/pot, thinned to 3 plants/pot Planting date: November 15, 2010
Experimental design: A dilution series of Vitazyme was prepared and applied to pots in a replicated greenhouse setting, the purpose of which was to determine relative degrees of response to the active agents at varying concentrations. A completely randomized design was used, with eight replications.

Treatment	Vitazyme concentration
	%
1	0
2	0.00001 (10 ⁻⁵)
3	0.0001 (10 ⁻⁴)
4	0.001 (10 ⁻³)
5	0.01 (10 ⁻²)
6	0.1 (10 ⁻¹)
7	1 (10 ⁰)
8	10 (10 ¹)
9	100 (10 ²)

Fertilization: none

Vitazyme application: At planting, 100 ml of the eight dilutions were added to the soil surface of the appropriate pots, directly after seeding.


Plant height results: Plant heights were measured for the three plants of each pot at harvest, on December 15, 2010, and averaged.



Treatment P	0.0000***
Model P	0.0000***
CV	7.55%
LSD _{0.05}	5.2 cm

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.

Dry weight results:



Treatment P	0.0000***
Model P	0.0000***
CV	7.55%
LSD _{0.05}	5.2 cm

¹Means followed by the same letter are not significantly different at P=0.05 according to the Student-Newman-Keuls Test.

Conclusions: Vitazyme at all concentrations improved the growth of corn above the control in terms of plant height and dry weight. All concentrations, from 10⁻⁵ to 10², gave significant responses for plant height, with the 10⁻² and 10⁻¹ dilutions giving the strongest responses (28% and 29%, respectively). Plant weight increases were statistically equal across the entire range of concentrations except for 10⁻⁵ and 10⁰%, the highest values appearing at 10⁻⁴, 10⁻³, and 10¹%; the reason for this bimodal response is not known. The field application rates are typically represented by 10⁻² to 10⁻³% Vitazyme.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher/Farmer: Nguyen Ngoc Tuan
Province, Viet Nam

Location: Easling Town, Cu Jut District, Dak Nong
Variety: 30 D55

Planting season: Summer – Fall, 2011

Experimental design: A corn field was divided into a Vitazyme treated area of 0.5 ha, and an untreated control area of 0.3 ha, to determine the effect of this product on the yield and profitability of the crop.

1. Control

2. Vitazyme

Vitazyme application: (1) 5% Vitazyme seed spray just before planting; (2) 1 liter/ha sprayed on the leaves and soil 35 days after planting


Growth results: With Vitazyme the following effects were noted:

- Taller plants
- Stronger root systems
- Darker green leaves
- Excellent disease resistance from steaked leaf and stem rot
- During a drought period, plants showed no leaf rolling
- Seed color was brighter

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	8.0	—
Vitazyme	9.0	1.0 (+13%)

**Increase in corn yield with
Vitazyme: 13%**



Income results:

Treatment	Vitazyme ¹ VND/ha	Total costs VND/ha	Total income ² VND/ha	Net income VND/ha	Extra profit VND/ha
Control	0	13,500,000	36,000,000	22,500,000	—
Vitazyme	500,000	14,000,000	38,250,000	26,500,000	4,000,000

¹VND = Vietnamese dollar; 1 USD = 20,000 VND.

²Corn price = 4,500 VND/kg.

**Increase in income with Vitazyme:
4,000,000 VND, or \$200.00/ha**

Conclusion: This Vietnamese corn trial revealed that Vitazyme, applied on the seeds and also 30 days after planting, increased yield by 13%, while boosting profit by \$200.00/ha. This program is highly effective for corn production in Viet Nam.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2011 Crop Results

Vitazyme on Corn

Researcher: Unknown

Research organization: National Academy of Agrarian Sciences, Vinnytsia State Agricultural Research Station

Location: Vinnytsia, Ukraine (Central Forest and Steppe Region)

Variety: Sangriya

Planting date: unknown

Soil type: gray podzolic (organic matter = 2.2%, hydrolyzed N = 8.4 mg/100 g soil, P = 15.8 mg/100g soil, exchangeable K = 12.4 mg/100 g soil, pH = 5.5)

Experimental design: Replicated corn plots were prepared and treated with a Vitazyme treatment, and plots treated with Vitazyme were also used to evaluate a carryover effect, to evaluate the effect of the product on corn yield and profitability.

1. Control


2. Vitazyme on leaves and soil

3. Vitazyme on leaves and soil in 2010

Vitazyme applications: 1 liter/ha at the 7 to 8-leaf stage (June 16, 2011) for Treatment 2

Yield results:

Treatment	Yield tons/ha	Yield change tons/ha
Control	8.9	—
Vitazyme foliar	10.2	1.3 (+15%)
Vitazyme in 2010	9.4	0.5 (+6%)



Yield increase with a Vitazyme foliar treatment: 15%

Yield increase with a Vitazyme in 2010: 6%

Income results:

- Income increase with a Vitazyme treatment: 2,050 hrn/ha
- Income increase with Vitazyme in 2010: 850 hrn/ha

Conclusion: This replicated corn trial in Vinnytsia, Ukraine, reveals what previous years' trials have shown ... that Vitazyme increases yield (+15% in 2011) and profitability (+2,050 hrn/ha in 2011) consistently. It is a product that powerfully improves agricultural productivity in Ukraine.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2009 Crop Results

Vitazyme on Corn

Researcher: Nathan Temples

Farm cooperater: Parker Brothers

Location: Sikeston, Missouri

Variety: Pioneer 33N58

Soil type: sandy loam

Planting rate: 31,500 seeds /acre

Row-spacing: 38 inches

Irrigation: unknown

Planting date: April 23, 2009

Experimental design: A corn field received Vitazyme on the seeds of 24 rows within the field, to determine the effects of this product on corn yield.

1. Control

2. Vitazyme

Fertilization: 200-60-90 lb/acre N-P₂O₅-K₂O


Vitazyme application: 8 oz/acre on the seeds at planting

Harvest date: September 23, 2009

Yield results:

Treatment	Yield -----bu/acre-----	Yield change
Control	193	—
Vitazyme	199	6 (3%)

**Increase in corn yield with
Vitazyme: 3%**



Conclusions: This high yielding corn study in Missouri revealed that Vitazyme, applied at 8 oz/acre to the seeds at planting, increased yield by 6 bu/acre (+3%). This is a highly profitable additional yield, to reveal the potential of Vitazyme to improve corn yields in the Corn Belt of Missouri.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2009 Crop Results

Vitazyme on Corn

Researcher: O.V. Kornijchuk, V.V. Plotnikov, and agronomic scientists

Organization: Vinnytsia State Agricultural Experiment Station, Ukraine Academy of Agrarian Sciences, Vinnytsia, Ukraine

Location: Ukraine central forest-steppe area near Vinnytsia

Planting date: May 22, 2009

Variety: Ronaldinio

Seeding rate: 22 kg/ha

Tillage: plowing, harrowing, and cultivation

Previous crop: winter wheat

Soil type: gray forest steppe soil; in the 0-30 cm layer, 2.2% organic matter, 8.4 mg/100 g of soil “hydrolyzed nitrogen”, 15.8 mg/100g of soil phosphorous, 12.4 mg/100 g of soil exchangeable potassium, and pH=5.5.

Experimental design: A uniform field was divided into plots of 1.0 ha each with two treatments and four replications. The objective of the study was to evaluate the effect of Vitazyme as either a seed application, or a seed plus foliar application, on the yield of corn grain.

1. Control

2. Vitazyme, once foliar


Fertilization: 60 kg/ha, 30 kg/ha P₂O₅, and 60 kg/ha K₂O

Vitazyme application: Treatment 2 received 1.0 liter applied to the leaves and soil on June 25, 2009, at the 7 to 8 leaf stage

Yield results:

Treatment	Grain yield tons/ha	Change tons/ha
Control	8.3	—
Vitazyme	9.0	0.7 (+8%)

**Increase in corn yield with
Vitazyme: 8%**



Income results:

Income increase with Vitazyme foliar: 479 hrn/ha

Conclusions: Corn grain with and without Vitazyme (1 liter/ha, foliar) in this Ukraine study showed an 8% yield increase. Moreover, the return to the farmer was improved substantially.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2009 Crop Results

Vitazyme on Corn

Researcher: Nathan Temples

Farm cooperator: Schlosser Farms

Location: Perkins, Missouri

Variety: Pioneer 33 N 58

Soil type: silt loam

Planting date: April 20, 2009

Planting rate: 25,000 seeds /acre

Irrigation: none

Experimental design: Five acres of a 90-acre corn field were treated with Vitazyme, applied with the herbicide, to determine the product's ability to improve crop yields.

1. Control

2. Vitazyme

Fertilization: 160-50-90 lb/acre N-P₂O₅-K₂O preplant


Vitazyme application: 13 oz/acre with the herbicide, 42 days after planting

Harvest date: October 3, 2009

Yield results:

Treatment	Yield	Yield change
	-----bu/acre-----	
Control	201	—
Vitazyme	206	5 (2.5%)

**Increase in yield with Vitazyme:
2.5%**



Conclusions: A Missouri corn study showed that Vitazyme, applied along with a herbicide at 13 oz/acre, increased yield by 5 bu/acre (+2.5%)

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2009 Crop Results

Vitazyme on Corn

Researcher: Nathan Temples

Farm cooperater: Seyer Farms

Location: Oran, Missouri

Variety: Dekalb

Soil type: sandy

Planting rate: 29,000 seeds /acre

Row-spacing: 30 inches

Irrigation: furrow, six times

Planting date: April 22, 2009

Experimental design: An 80-acre irrigated corn field was divided into 60 acres treated with Vitazyme, and 20 acres left untreated, to determine the product's effect on crop yield.

1. Control

2. Vitazyme

Fertilization: unknown


Vitazyme application: 13 oz/acre along with a herbicide

Harvest date: September 14, 2009

Yield results:

Treatment	Yield	Yield change
	----- bu/acre -----	
Control	210	—
Vitazyme	217	7 (3.3%)

**Increase in corn yield with
Vitazyme: 3.3%**



Conclusions: In this high yielding Missouri irrigated corn trial, Vitazyme increased corn yield by 7 bu/acre (3.3%), showing the utility of this product to improve production even at high yield levels.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2009 Crop Results

Vitazyme on Corn

Researcher: Nathan Temples

Farm cooperators: Donnie and Chris Wondel, D and C Farms

Location: Arbor, Missouri

Variety: 32 D 78

Soil type: silt loam

Planting rate: 29,000 seeds /acre

Row-spacing: 30 inches

Irrigation: furrow, six times

Planting date: May 7, 2009

Experimental design: A 55-acre field was treated with Vitazyme on 40 acres, using seed and foliar treatments, to determine if this product would increase the yield of grain.

1. Control

2. Vitazyme

Fertilization: unknown


Vitazyme application: (1) 8 oz/acre on the seeds at planting; (2) 13 oz/acre on the leaves and soil at 7 inches height, with a herbicide

Harvest date: October 6, 2009

Yield results:

Treatment	Yield -----bu/acre-----	Yield change
Control	175	—
Vitazyme	190	15 (9%)

Increase in yield with Vitazyme: 9%



Conclusions: In this Missouri corn trial, using seed and foliar/soil applications, the Vitazyme treatments increased grain yield by 9% (15 bu/acre) in this high-yielding field, showing the program's great effectiveness in corn programs, even when yields are high.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2008 Crop Results

Vitazyme on Corn

A Long-Term study: Year 1

Researcher: Bertel Schou, Ph.D. Research organization: ACRES (Agricultural Custom Research and Environmental Services), Cedar Falls, Iowa

Variety: Pioneer 34R67 (BBCH Scale:BCOR)

Planting rate: 29,900 seeds/acre

Planting date: May 21, 2008

Tillage: conventional (cultivated and harrowed on May 21)

Row spacing: 30 inches

Planting depth: 2 inches

Previous crop: corn

Soil type: Kenyon loam (34% sand, 46% silt, 20% clay, 3.6% organic matter, 15.0 meq/100 grams cation exchange capacity, pH 7.2, good fertility)

Soil test results, initial for all plots (analyzed May 15, 2008): pH, 7.2; organic matter, 3.90%; N, 89 lb/acre; SO₄-S, 6 lb/acre; P₂O₅, 1,076 lb/acre; Ca, 5,407 lb/acre; Mg, 916 lb/acre; K, 298 lb/acre; Na, 52 lb/acre; B, 1.76 lb/acre; Fe, 460 lb/acre; Mn, 176 lb/acre; Cu, 3.4 lb/acre; Zn, 12.6 lb/acre; base saturations: Ca, 72.6%, Mg, 20.5%, K, 2.1%, Na, 0.6%, others, 4.2%

Experimental design: A field was selected to place plots (15 x 40 feet) in a randomized complete block design (five replicates), using two treatments for a long-term field study. These plots are designed to assess the long-term effects of Vitazyme on the yield and growth of corn and soybeans in rotation, but moreover the effects on the physical, chemical, and microbial characteristics of the soil.

1. Control

2. Vitazyme

Fertilization: 120 lb/acre of N as 28% N applied postemergence in 20-inch spaced bands, using drop nozzles from a shielded sprayer

Vitazyme application: 13 oz/acre (1 liter/ha) in the seed furrow at planting (May 21), and 13 oz/acre sprayed over the leaves and soil on July 6, 2008, at the V6 stage

Weed control: Harness Extra preemergent, and Accent postemergent

Microorganism analyses: Soil biological activity was evaluated in the spring and fall to determine product effects on a number of parameters. Soil samples were collected from the root zones of plants from each of the five replicates, and then combined for each treatment and sent to the Soilfoodweb Laboratory in Corvallis, Oregon, for analysis.

May 29 analysis (baseline values for future comparisons)

Treatment	Organism biomass	Active bacteria	Total bacteria	Active fungi	Total fungi	Protozoa			Total nematodes
						Flagellates	Amoebae	Ciliates	
Control	0.81	28.4	1,853	20.4	244	5,718	17,211	34	2.33
Vitazyme	0.80	24.7	2,324	13.3	282	3,454	5,738	72	1.84

Treatment	Total fungi to bacteria	Active fungi to total fungi	Active bacteria to total bacteria	Active fungi to active bacteria	Plant-available nitrogen
					lb/acre
Control	0.13	0.08	0.02	0.72	75 to 100
Vitazyme	0.12	0.05	0.01	0.54	50 to 75


September 10 analysis

Treatment	Organism biomass	Active bacteria	Total bacteria	Active fungi	Total fungi	Protozoa			Total nematodes
						Flagellates	Amoebae	Ciliates	
		ug/g	ug/g	ug/g	ug/g	no./g	no./g	no./g	no./g
Control	0.82	41.6	929	18.7	352	1,690	5,618	169	2.73
Vitazyme	0.81	3.4	1,033	13.4	240	8,594	7,103	103	0.57


Treatment	VA mycorrhizae	Total fungi to bacteria	Active fungi to total fungi	Active bacteria to total bacteria	Active fungi to active bacteria	Plant-available nitrogen
	% infection					lb/acre
Control	0	0.38	0.05	0.04	0.45	50 to 75
Vitazyme	0	0.23	0.06	0.03	0.41	75 to 100

Differences in microbes between the two treatments are not pronounced, although there was a distinctly greater number of protozoa with the Vitazyme treatment, especially flagellates. There was no VAM mycorrhizal root infection for either treatment. Fungal and bacterial ratios were not very different, but plant-available nitrogen was decidedly greater with the Vitazyme treatment.

Total Protozoa



Plant-Available Nitrogen



Increase in protozoa with Vitazyme: 111%


Increase in N availability with Vitazyme: 25 lb/acre

Harvest date: The crop was harvested on November 1, 2008, using a Massy-Ferguson 8 plot combine. Two rows 40 feet long were harvested from each plot.

Plant populations: The populations of the two treatments were very similar: 21,414 plants/acre for the control, and 21,235 plants/acre for the Vitazyme treatment. This difference was not significant (P=0.871).

Yield results:

Treatment	Grain yield	Yield increase
	bu/acre	bu/acre
Control	114.96 b	—
Vitazyme	127.02 a	12.06 (+10%)
LSD (0.05)	8.53	
Standard deviation	4.86	
Coeff. of variation	4.02	
Replicate F	4.4	
Replicate probability	0.090	
Treatment F	15.4	
Treatment probability	0.017	



Vitazyme significantly increased the yield of corn, by 12.06 bu/acre, a full 10% above the control yield.

Conclusions: In this first year of a long-term trial to evaluate the effects of Vitazyme on the physical, chemical, and microbiological effects of the soil, and on crop response, Vitazyme greatly boosted grain yield (12.06 bu/acre, or 10%) above the control. Baseline soil chemical analyses were completed, as were baseline microbiological analyses. A September 10 microbial analysis revealed that, while both treatments showed minimal differences in most parameters measured, there was a marked 111% increase in total protozoa with Vitazyme. In addition, the supply of plant-available nitrogen was improved by about 25 lb/acre with Vitazyme, a significant factor in the current climate of high and volatile fertilizer prices. Work will continue during the coming years on monitoring the changes brought about by Vitazyme on an array of soil and plant characteristics.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2008 Crop Results

Vitazyme on Corn

Researcher/Farmer: Rick Nichols

Variety: Pioneer 34Y88 (non-GMO)

Row spacing: 30 inches

Planting date: May 4, 2008

Experimental design: A field was divided into a control area receiving no sidedressed nitrogen or Vitazyme, and a treated area receiving both. The objective of the test was to evaluate the effect of combined sidedressed nitrogen plus Vitazyme on crop yield.

Location: Hebron, Indiana

Soil type: silty clay "gumbo"

Population: 34,000 seeds/acre

Previous crop: soybeans

1. Control

Fertilization: Before planting: 140 lb/acre nitrogen, as urea. At planting: 300 lb/acre 18-46-60% N-P₂O₅-K₂O placed 4 inches to the side of the seeds. At sidedressing, in June (corn about 2 feet tall): 40 lb/acre nitrogen as a 28% solution

Vitazyme application: 13 oz/acre with sidedressed nitrogen on the treated area, applied in June at the 2-foot height


Harvest date: October 7, 2008

Yield results: Six rows of field length were harvested and weighed from each treatment in passes near one another. However, no row length was measured, so per acre yields were not obtained.

2. Vitazyme + sidedressed nitrogen

Treatment	Yield	Increase
	bu/6 rows	bu/6 rows
1. Control	120	—
2. Vitazyme + Sidedressed N	140	20 (+17%)

Increase in corn yield: 17%



Conclusions: In this northern Indiana corn trial, Vitazyme side-dressed with 40 lb/acre of nitrogen as a 28% solution increased the yield by 17% above the control. It was not possible to separate the effects of the nitrogen and the Vitazyme, but it is well documented that Vitazyme enhances the utilization of soil and fertilizer applied nutrients, especially nitrogen.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2008 Crop Results

Vitazyme on Corn

Researcher/Farmer: Gary Burkey

Variety: Flexseed 4918 non-GMO

Row spacing: 30 inches

Planting date: May 6, 2008

Soil test: pH, 6.6; cation exchange capacity, 24.6 meq/100g; organic matter, 3.6%; base saturations, Ca = 65.6%, Mg = 21.8%, K = 1.4%, Na = 0.3%, other bases = 4.8%, H = 6.0%; estimated N release, 86 lb/acre; S, 9 ppm; P₂O₅, 175 lb/acre; Ca, 6,468 lb/acre; Mg, 1,290 lb/acre; K, 274 lb/acre; Na, 38 lb/acre; B, 0.9 ppm; Fe, 1,842 ppm; Mn, 77 ppm; Cu, 1.1 ppm; Zn, 26.7 ppm

Experimental design: A corn field was treated entirely on the seeds with Vitazyme, and part of the field received a foliar Vitazyme treatment as well, along with two other products in the sprayer tank. The objective of the study was to evaluate the effect of an additional Vitazyme application and these other foliar products on corn yield.

1. Vitazyme on the seeds

2. Vitazyme on the seeds, plus Vitazyme and two other products on the leaves

Fertilization: *Before planting:* 150 lb/acre potassium chloride (0-0-60% N-P₂O₅-K₂O); 100 lb/acre diammonium phosphate (18-46-0% N-P₂O₅-K₂O); 70 lb/acre N from dry urea. *At planting:* 4 gallons/acre 3-18-18% N-P₂O₅-K₂O on the seeds. *At knee-height:* 70 lb/acre N (28% N) side-dressed with a row-crop cultivator. *Foliar spray on July 6:* Tricert K (1 quart/acre of a 50-0-20 N-P₂O₅-K₂O material), manganese (1.5 lb/acre) with Vitazyme.

Vitazyme application: (1) 13 oz/acre on the seeds at planting, along with 3-18-18 fertilizer; (2) 13 oz/acre sprayed foliar with Tricert K and manganese on July 6


Weather results: a wet spring and late planting, few rains in July, and a very dry late July and August, followed by a 12-inch flooding rain in mid-October

Harvest date: November 10, 2008

Yield results: Eight -row swatches were combined and weighed for both treatments. Due to an extreme rain event in mid-October, water rose so high in the field that the ears were covered for two to three days. In spite of that problem the corn grade was not affected, although untreated corn from neighbors' fields suffered water damage to their grain.

Treatment	Yield bu/acre	Change bu/acre
1. Vitazyme on seeds	204	—
2. Vitazyme on seeds + foliar with Tricert K + Mn	241	37 (+18%)

Increase in corn yield: 18%



Conclusions: In this Indiana in-field corn trial, Vitazyme plus Tricert K and manganese boosted the yield by 18% (37 bu/acre), though it was not possible to separate the individual effects of these products. Vitazyme works in synergism with native soil and applied nutrients to boost utilization, so this great yield increase is not uncommon. **Of great interest is the fact that submersion of the ears before harvest for up to three days did not reduce the grain quality, indicating that cell wall integrity and anti-pathogen properties of the grain were likely enhanced by Vitazyme throughout the field.**

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2008 Crop Results

Vitazyme on Corn

Researcher/Farmer: Gary Burkey

Variety: Flexseed 303 Triple-Stack

Row spacing: 30 inches

Planting date: May 18, 2008

Soil test: pH, 7.2; cation exchange capacity, 22.53 meq/100g; organic matter, 6.0%; base saturations, Ca = 70.8%, Mg = 20.5%, K = 4.2%, Na = 0.3%, others = 4.2%, H = 0%; estimated N release, 105 lb/acre; S, 10 ppm; P₂O₅, 468 lb/acre; Ca, 6,376 lb/acre; Mg, 1,110 lb/acre; K, 739 lb/acre; Na, 31 lb/acre; B, 0.9 ppm; Fe, 277 ppm; Mn, 29 ppm; Cu, 0.4 ppm; Zn, 6.6 ppm

Experimental design: A field was treated entirely with Vitazyme on the seeds at planting, and a portion of the field was foliar treated to determine the effect of this later application on crop yield.

1. Vitazyme on the seeds

Location: Coats, Indiana

Soil type: mucky sand

Population: 31,000 seeds/acre

Soil test: pH, 7.2; cation exchange capacity, 22.53

meq/100g; organic matter, 6.0%; base saturations, Ca = 70.8%, Mg = 20.5%, K = 4.2%, Na = 0.3%, others = 4.2%, H = 0%; estimated N release, 105 lb/acre; S, 10 ppm; P₂O₅, 468 lb/acre; Ca, 6,376 lb/acre; Mg, 1,110 lb/acre; K, 739 lb/acre; Na, 31 lb/acre; B, 0.9 ppm; Fe, 277 ppm; Mn, 29 ppm; Cu, 0.4 ppm; Zn, 6.6 ppm

Experimental design: A field was treated entirely with Vitazyme on the seeds at planting, and a portion of the field was foliar treated to determine the effect of this later application on crop yield.

2. Vitazyme on the seeds + leaves

Fertilization: Before planting: 150 lb/acre potassium chloride (0-0-60% N-P₂O₅-K₂O); 100 lb/acre diammonium phosphate (18-46-0% N-P₂O₅-K₂O). At planting: 4 gallons/acre 3-18-18% N-P₂O₅-K₂O on the seeds at planting. Sidedressed on June 18, at 5-foot plant height: 40 gallons/acre 28% N.

Vitazyme application: (1) 13 oz/acre on the seeds at planting, with 3-18-18% N-P₂O₅-K₂O over all areas; (2) 13 oz/acre foliar over one portion of the field, on June 18


Weather results: a wet spring and late planting, few rains in July, and very dry in late July and August, with a flooding rain (12 inches) in mid-October

Harvest date: December 12, 2008

Yield results: Eight-row swaths were combined and weighed for both treatments.

Treatment	Yield bu/acre	Change bu/acre
1. Vitazyme on seeds	155	—
2. Vitazyme on seeds + leaves	160	5 (+3%)

Conclusions: Vitazyme applied foliar in this northern Indiana corn trial resulted in a substantial 5 bu/acre increase in yield above the treatment receiving only a seed treatment.



Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2008 Crop Results

Vitazyme on Corn

A Greenhouse Study

Researcher: Paul W. Syltie, Ph.D.
Texas

Location: Vital Earth Resources Research Greenhouse, Gladewater,
Variety: yellow dent

Planting date: January 31, 2008

Soil type: silt loam

Pot size: 1 gallon

Planting rate: 10 seeds/pot, thinned to three plants

Watering: on-demand

Temperature: 55 to 85°F

Planting depth: 0.5 inch

Experimental design: A replicated greenhouse pot study was designed to evaluate the effect of various rates of Vitazyme on the growth of corn. six replicates were included with six treatments, and the data were analyzed using Analysis of Variance with CoHort software.

1. Control

4. Vitazyme at 26.0 oz/acre

2. Vitazyme at 7.5 oz/acre

5. Vitazyme at 39.0 oz/acre

3. Vitazyme at 13.0 oz/acre

6. Vitazyme at 52.0 oz/acre

Vitazyme applications: The 13.0 oz/acre application was made immediately after planting to the soil surface of the pot, using 100 ml of a 0.0016% solution. Other treatments were multiples of this rate.

Harvest date: On March 5, 35 days after planting, the soil was washed from the roots of the plants, and measurements were made of the height of each plant. The plants were then placed in a drying oven at about 50°C for 48 hours.

Plant height results:

Treatment	Plant height	Height change*
	cm	cm
5 (Vitazyme, 3x)	85.3 a	5.5 (+7%)
3 Vitazyme, 1x)	84.6 a	4.8 (+6%)
4 (Vitazyme, 2x)	83.8 a	4.0 (+5%)
2 (Vitazyme, 0.5x)	83.4 a	3.6 (+5%)
6 (Vitazyme, 4x)	80.0 b	0.2 (+0%)
1 (Control)	79.8 b	—
Statistical analysis		
Replicate P	0.0245	
Treatment P	0.0040	
Model P	0.0029	
Coefficient of variation	3.24%	
LSD _{0.10}	2.6 cm	

*Compared to the untreated control, Treatment 1.

The highest (4 times normal) rate, as well as the untreated control, gave significantly shorter plants than all of the other Vitazyme treatments. There was no statistical difference among the 0.5 to 3 times normal treatments, but the greatest heights were for the 3 times normal and normal treatments.

Increase in plant height

3x Vitazyme	7%
1x Vitazyme	6%
2x Vitazyme	5%
0.5x Vitazyme	5%

Dry weight results:


Treatment	Dry weight	Dry weight change*
	cm	cm
2 (Vitazyme, 0.5x)	9.30 a	1.11 (+14%)
4 Vitazyme, 2x)	9.30 a	1.11 (+14%)
6 (Vitazyme, 4x)	9.22 a	1.03 (+13%)
5 (Vitazyme, 3x)	9.01 a	0.82 (+10%)
3 (Vitazyme, 1x)	9.00 a	0.81 (+10%)
1 (Control)	8.19 b	—

Statistical analysis	
Replicate P	0.0092
Treatment P	0.0192
Model P	0.0043
Coefficient of variation	6.23%
LSD _{0.10}	0.55 gram

*Compared to the untreated control, Treatment 1.

All of the Vitazyme applications gave increases in dry plant weight of from 10 to 14%, the highest increases being with the 0.5x, 2x, and 4x rates. None of these differences were statistically significant, and all exceeded the control.

Increase in plant dry weight	
0.5x Vitazyme	14%
2x Vitazyme	14%
4x Vitazyme	13%
3x Vitazyme	10%
1x Vitazyme	10%



Conclusions: In this greenhouse study to evaluate the effects of progressively higher rates of Vitazyme to stimulate corn height and dry weight accumulation, the product proved to significantly increase plant height by 5 to 7% at all but the 4x (52 oz/acre) rate, whereas dry weight significantly increased from 10 to 14% for all of the Vitazyme rates. These data prove that more than just the standard 13 oz/acre rate can be effective in stimulating crop growth, but higher rates do not produce a linear yield or growth increase.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2007 Crop Results

Vitazyme on Corn

Researchers: Fred Vaughn and Greg Wilson

Organization: Vaughn Agricultural Research Services

Location: Branchton, Ontario, Canada

BBCH Scale: BCOR

Variety: Pioneer 38P03

Planting rate: 76,000 seeds/ha

Planting depth: 5cm

Row spacing: 76 cm

Planting Date: May 14, 2007

Seedbed conditions: dry, fine

Soil temperature at planting: 13.3C
meq/100 g CEC, good fertility

Soil: silt loam (31.9% sand, 53.7% silt, 14.4% clay), 6.2 pH, 14.2

Field preparation: cultivation twice

Previous crop: 2004, winter wheat (with Cobutox); 2005, potatoes (with Dual + Sencor); 2006, corn (Dual + Marksman)

Experimental design: A uniform site was divided into plots that were 3x6 meters (six rows), using four treatments with six replications in a randomized complete block design. The objective of the study was to determine Vitazyme's ability to improve soybean yield with two applications. The treatments were as follows:

Treatment	At planting	At 8 leaves	Nitrogen
	----- liters/ha -----		Kg/ha
1. No Vitazyme	0	0	60
2. No Vitazyme	0	0	60
3. Vitazyme	1	1	120
4. Vitazyme	1	1	120

Fertilization: All areas received 200 kg/ha of dry 6-24-24% N-P₂O₅-K₂O before planting. 100 liters/ha of liquid 6-24-6% N-P₂O₅-K₂O was applied in the seed furrow at planting (May 14). A 28% nitrogen solution was applied to the plots on June 8 so that the appropriate plots would receive either 60 or 120 kg/ha of nitrogen.

Vitazyme application: To Treatments 3 and 4, 1 liter/ha was applied to the seeds at planting (May 14), as a spray on the seeds just behind the disc openers, and 1 liter/ha was applied to the leaves and soil at the eight-leaf stage (June 20).

Crop emergence date: May 18, four days after planting


Weed control: unknown

Harvest date: October 12, 2007. An area of 1.52 x 6.00 meters (the two center rows) was harvested for each plot.

Yield results: There were no significant differences in grain moisture content and test weight, nor were any differences discovered in stalk lodging. Thus, those data are not included below.

Grain Yield

Treatment	Grain yield	Change*
	bu/acre	bu/acre
1. No Vitazyme, 60 N	118.6 c	—
2. No Vitazyme, 120 N	144.3 b	25.7 (+22%)
3. Vitazyme, 60 N	137.7 b	19.1 (+16%)
4. Vitazyme, 120 N	166.8 a	48.2 (+41%)
LSD	16.3	
CV	11.37	
Bartlett's X2	3	
P (Bartlett's X2)	0.392	
Replicate F	4.315	
Replicate Prob (F)	0.0124	
Treatment F	9.158	
Treatment Prob (F)	0.0011	



Vitazyme Effect at 60 kg/ha N

Treatment	Yield	Change
	bu/acre	bu/acre
No Vitazyme	118.6 b	—
Vitazyme	137.7 a	19.1 (+16%)

Yield increase with Vitazyme at 60 kg/ha N: 16%

Vitazyme Effect at 120 kg/ha N

Treatment	Yield	Change
	bu/acre	bu/acre
No Vitazyme	144.3 b	—
Vitazyme	166.8 a	22.5 (+16%)

Yield increase with Vitazyme at 120 kg/ha N: 16%

At both nitrogen levels, Vitazyme significantly increased grain yield at P=0.10. This increase was 16% above the control at both nitrogen levels. What is especially interesting to note is that the 60 kg/ha N yield (137.7 bu/acre) with Vitazyme was statistically equal to the 120 kg/ha N yield (144.3 bu/acre) without Vitazyme. This reveals a benefit of Vitazyme to improve the utilization of fertilizer nitrogen.

Income results: At \$4.00/bu, the increased incomes for the grain produced in this study are as follows:

At 60 kg/ha N. No Vitazyme: 118.6 bu/acre x \$4.00/bu = \$474.40

Vitazyme: 137.7 bu/acre x \$4.00/bu = \$550.80

Increase with Vitazyme: \$74.40/acre

At 120 kg/ha N. No Vitazyme: 144.3 bu/acre x \$4.00/bu = \$577.20

Vitazyme: 166.8 bu/acre x \$4.00/bu = 667.20

Increase with Vitazyme: \$90.00/acre

Conclusions: In this southern Ontario, Canada, study of Vitazyme on corn at two nitrogen levels, Vitazyme was shown to significantly increase grain yield, by 16% above the respective control (no Vitazyme) levels. Moreover, the yield of the Vitazyme + 60 kg/ha N rate was statistically equal to the 120 kg/ha N rate without Vitazyme, demonstrating the ability of the product to improve the utilization of nitrogen. Two applications of 1 l/ha, at planting and again at the eight-leaf stage, brought about this yield improvement. The yield increases gave significant income increases: \$74.40/acre at 60 kg/ha nitrogen, and \$90.00/acre at 120 kg/ha nitrogen.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2006 Crop Results

Vitazyme on Corn

Ministry of Sugar, Cuban Ministry of Agriculture

Researchers: Wilberto G. Marrero and Jorge G. Acosta

Location: Juan Abrahantes Farm, Madruga, Havana Province, Cuba

Variety: unknown

Soil type: red ferralitic of low fertility

Planting rate: unknown

Row spacing: unknown

Planting date: July 23, 2006

Watering: rain-fed

Experimental design: An area of 1.5 acres in a production corn field was treated with Vitazyme twice, each time at 1 liter/ha, to determine the effect of the product on corn yield.

1. Control

2. Vitazyme

Fertilization: unknown

Vitazyme application: 1 liter/ha on the leaves and soil on August 7, 15 days after planting, and again 39 days later on September 15

Harvest date: October 14, 2006, 83 days after planting


Growth results:

Parameter	Control	Vitazyme
Rows of kernels	Averaging <12 rows/ear	Averaging >12 rows/each
Ear size	Average size	Larger than average
Stalk diameter	Average diameter	Greater than average diameter
Plant vigor	Average vigor	More vigorous
Plant height	2.00 meters average	1.55 meters average
Root development	Moderate	Extensive

Yield results:

Treatment	Yield tons/ha	Increase tons/ha
Control	1.45	—
Vitazyme	2.80	1.35 (+93%)
Historical yield	0.70	

Increase in corn yield: 93%



Conclusions: This Cuban corn study showed that Vitazyme greatly increased corn yield (by 93%) with two applications at 1 liter/ha each time, separated by 39 days. **This yield was four times the normal historical yield experienced in that area under the management system used.** The treated corn plants expressed superior vigor and growth throughout the growth cycle.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2006 Crop Results

Vitazyme on Corn

Researchers: Eng. Wilberto Gonzalez, and Eng. Jorge Gonzalez, Camilo Cienfuegos, Agricultural Enterprise

Location: Armistad Farm of Camilo Cienfuegos Agricultural Enterprise, Havana Province, Cuba

Variety: unknown

Soil type: red ferralitic

Planting date: late 2005 to early 2006


Experimental design: A commercial production trial involved a split field area of 1.0 ha treated and 1.0 ha untreated with Vitazyme at Armistad Farm.

1. Control

2. Vitazyme

Fertilization: unknown

Vitazyme applications: 1.0 liter/ha on the leaves twice, separated by 30 days



Increase in corn yield: 415%

Conclusions: This commercial corn trial in Cuba revealed the remarkable ability of Vitazyme to increase corn production, with a very large 415% yield increase.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2006 Crop Results

Vitazyme on Corn

USDA/National Soil Tilth Laboratory

Researcher: Jerry Hatfield, Ph.D.

Variety: Pioneer 35P17

Planting date: April 18, 2006

Tillage: chisel plowing on November 26, 2005, and field cultivation on April 4 and April 18, 2006

Experimental design: A field area was treated with Vitazyme to determine if corn planted in a double-row fashion would respond to the product under conventional tillage. The trial was non-replicated.

Location: Ames, Iowa

Planting rate: 36,000 seeds/acre in double rows

1. Control

Fertilization: 50 lb/acre of N as 32% UAN on October 18, 2005; 30-80-120 lb/acre of N-P₂O₅-K₂O dry spread on November 23, 2005; 300 lb/acre of SuperCal 98 pelleted lime on February 8, 2006; 300 lb/acre of SuperCal SO₄ pelleted gypsum on February 9, 2006; sidedressed 180 lb/acre of N as 32% UAN on May 29, 2006

Vitazyme application: 13 oz/acre foliar, hand applied, on June 5, 2006; 13 oz/acre foliar, hand applied, on June 27, 2006


Herbicide application: 3 qt/acre of Lumax, pre-emergent, on April 24, 2006

Harvest date: October 19, 2006

Yield results:

Treatment	Yield	Increase
	tons/ha	tons/ha
Control	205	—
Vitazyme	215	10 (+5%)

**Yield increase with Vitazyme:
5%**



Conclusions: On this non-replicated corn-yield study in central Iowa, using a double row system, two foliar Vitazyme applications utilizing 13 oz/acre each time increased the grain yield by 10 bu/acre (5%). Had a seed application been made it is likely that the response would have been greater.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2006 Crop Results

Vitazyme on Corn (Organic)

Research location: Ontario County, New York

Planting date: May 24

Row spacing: 30 inches

Experimental design: A field of organically grown corn was divided into Vitazyme treated and untreated areas in an effort to determine the product's effects on the yield of high-moisture corn. This corn was placed in an air-tight silo to be ground and used for cattle feed later.

Variety: Blue River 42A32 (96-day)

Soil type: silty clay loam

Seeding rate: 28,400/acre

1. Control

2. Vitazyme

Fertilization: Liquid cow manure, 8,500 gal/acre in November of 2005, and 7,300 gal/acre on May 8, 2006; 300 lb/acre of a 5-5-5% N-P₂O₅-K₂O dry Fertrell organic mix, in a 2 x 2-inch placement


Vitazyme treatment: 13 oz/acre on the seed in-furrow at planting

Weather for 2006: adequate moisture until August, then excessive afterwords

Harvest date: October 13

Yield results: Each parcel was harvested for a certain area, and the grain was dumped into a bin where a measurement of volume was taken. From the difference of these values the value of each treatment was calculated based upon the average field yield.

Treatment	Grain yield tons/acre	Yield increase tons/acre
Control	3.39	—
Vitazyme	3.81	0.42 (+12%)



**Grain increase with Vitazyme:
12%**

Income results: The value of high moisture corn (25%) is about \$175/ton.

Value of control corn \$593.25/acre
Value of Vitazyme corn \$666.75/acre
Income increase with Vitazyme \$73.50/acre

Conclusions: In this New York split-field study on organically grown corn, only one 13 oz/acre treatment of Vitazyme, on the seeds, produced a marked 12% increase in the yield of high moisture corn. This yield increase translated into an additional \$73.50/acre income, or about an \$18 return for each dollar invested in Vitazyme.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2005 Crop Results

Vitazyme on Corn

North Carolina State University

Researcher: Ron Heiniger, Ph.D.

Variety: De Kalb Dk 69-71 RR/YG

Soil type: Roanoke silt loam

Row width: 30 inches

Experimental design: A split-plot randomized complete block design (four replicates) was placed on a uniform soil area with the main plots containing the two seeding rates. Plots were 10 x 40 feet. Subplots contained starter fertilizer, starter fertilizer + Vitazyme, Vitazyme only, and a control. Evaluations were made on stalk diameter, root parameters, and yield to discover the effects of all variables on these parameters.

Location: Hertford, North Carolina

Planting date: April 21, 2005

Previous crop: soybeans

Population: 27,000 and 38,000 seeds/acre

Main Plots

1. 27,000 seeds/acre
2. 38,000 seeds/acre

Subplots

1. Control
2. Vitazyme
3. Starter
4. Vitazyme + Starter


Fertilization: A 19-19-0% N-P₂O₅-K₂O fertilizer was applied to the subplots 3 and 4 at a 10 gal/acre rate in a 2 x 2 inch band below and beside the seeds at planting. On June 7, 60 gal/acre of 30% UAN (urea ammonium nitrate) was applied.

Vitazyme application: 13 oz/acre on the seeds at planting for subplots 2 and 4

Weed control: excellent control with Bicep, Roundup, and atrazine herbicides


Root and stalk results: In early July five plants in consecutive order in rows of each treatment were dug, and the soil was washed from the root balls. Roots were pruned and dried, and the stalk diameter at the first internode below each ear was measured. Root ball depth and diameter were also measured.

Root Ball Diameter




*Significantly greater than the control. $LSD_{0.05} = 1.32$ cm. (Comparisons are made within the same plant population.)


Root Mass



*Significantly greater than the control. $LSD_{0.05} = 43.5$ g



*Significantly greater than the control. $LSD_{0.05} = 0.93$ mm.



*Significantly greater than the control. $LSD_{0.05} = 22.9$ bu/acre (Comparisons are made within the same plant population.)

Conclusions (by the researcher): Significant treatment effects or interactions involving Vitazyme were found for the diameter of the root ball, root mass, stalk diameter and grain yield. In the case of the diameter of the root ball there was a significant plant population by treatment interaction. At the lower plant population of 28 000 plants/acre the combination of 19-19-0 and Vitazyme significantly increased the diameter of the root ball compared to either product used alone or when compared to the untreated check. Neither the 19-19-0 nor Vitazyme when used alone increased the diameter of the root ball compared to the untreated check. In contrast, at the higher plant population, Vitazyme, 19-19-0, or the combination of the two significantly increased the diameter of the root ball compared to the untreated check. Although none of these three treatments were significantly different from each other the combination of Vitazyme and 19-19-0 again tended to have the higher yield. There were no significant interactions for root mass. However, there was a significant treatment effect. Vitazyme when used alone or in combination with 19-19-0 resulted in greater root mass compared to the untreated check. Again, the combination of Vitazyme and 19-19-0 produced the greatest root mass when compared with either treatment used alone. There was also a treatment effect on stalk diameter. The combination of Vitazyme and 19-19-0 increased stalk diameter at the first internode below the ear when compared with the untreated check or with a treatment of only 19-19-0. There was not a significant difference in stalk diameter between a treatment with only Vitazyme and the combination of Vitazyme and 19-19-0. However, the combination did have the largest stalk diameter.

For grain yield there was a significant plant population and treatment interaction. At the lower plant population, Vitazyme alone significantly increased yield compared to either the starter fertilizer or the untreated check; while the combination of starter and Vitazyme resulted in a yield similar to that obtained by Vitazyme alone. In comparison, at the higher plant population, the starter treatment, Vitazyme, or the combination of the two resulted in statistically similar yields but only the Vitazyme or Vitazyme-starter combination had significantly higher yields than the untreated check.

In summary, Vitazyme or Vitazyme in combination with 19-19-0 increased root ball diameter, root mass, stalk diameter, and grain yield compared to an untreated check. While plant disease ratings were not taken in this study, it is unlikely that the Vitazyme effect was related to better disease resistance. It appears that Vitazyme applied to the seeds at planting improves early root development resulting in a larger root mass, greater stalk diameter, and increased yield.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2004 Crop Results

Vitazyme on Corn North Carolina State University

Researcher: Ron Heiniger, Ph.D

Location: Clarkton, North Carolina

Variety: DKC69-71 RRH62

Tillage: conventional

Population: 33,000 seeds/acre

Row width: 30 inches

Planting date: April 24, 2004

Previous crop: soybeans

Experimental design: A plot area of 18,000 ft² (180 x 100 ft) was divided into individual plots of 400 ft² (40 x 10 ft), with four replicates. The objective of the study was to evaluate the potential of Vitazyme bios-timulant to improve grain yield at five nitrogen rates.

Control

1. No N
2. 56 lb/acre
3. 112 lb/acre N
4. 224 lb/acre N
5. 280 lb/acre N

Vitazyme

6. No N
7. 56 lb/acre
8. 112 lb/acre N
9. 224 lb/acre N
10. 280 lb/acre N

Fertilization: 10 gal/acre of 19-19-0 %N-P₂O₅-K₂O in a 2x2 band on April 24 after broadcasting 30% UAN and a 10-34-0 fertilizer on April 19

Herbicide application: Lariat (3 qt/acre) on April 19, broadcast pre-plant

Insecticide application: Counter 20CR (7 lb/acre) on April 21, T-banded on April 21

Vitazyme application: 13 oz/acre on the seeds at planting, and 13 oz/acre broadcast at knee height

Harvest date: September 23, 2004

Yield Results:

Treatment	N-level	Yield	Change ¹	NCLB rating ²	Change ¹	GLS rating ³	Change ¹
	lb/acre	bu/acre	bu/acre				
Control	0	153.9	—	9.25	—	8.88	—
	56	170.2	—	9.38	—	8.88	—
	112	145.9	—	9.63	—	9.00	—
	224	159.3	—	9.75	—	9.38	—
	280	139.7	—	10.00	—	9.50	—
	Average		153.8	—	9.60	—	9.13
Vitazyme	0	181.6	27.7* (+18%)	9.38	0.13 (+1%)	8.88	0 (0%)
	56	162.6	(-) 7.6 (-4%)	9.75	0.37* (+4%)	9.25	0.37* (+4%)
	112	166.8	20.9* (+14%)	9.88	0.25* (+3%)	9.50	0.50* (+6%)
	224	178.6	19.3* (+12%)	9.88	0.13 (+1%)	9.63	0.25* (+3%)
	280	151.2	11.5 (+8%)	10.00	0 (0%)	9.50	0 (0%)
	Average		168.2	14.4* (+9%)	9.78	0.18 (+2%)	9.35
	LSD _{0.05}	14.3		0.19		0.15	


¹ Differences are compared with the same N level for control and Vitazyme

²NCLB = Northern Corn Leaf Blight: 1 = leaves covered with lesions, 10 = no disease.


³GLS = Gray Leaf Spot: 1 = leaves covered with lesions, 10 = no disease.

*Significant difference vs. the control at P=0.05.

Corn Grain Yield




Average of all plots




Vitazyme significantly boosted grain yield across all N levels, by 14.4 bu/acre (9%), and especially at the 0 N level (+18%), the 112 lb/acre N level (+14%), and the 224 lb/acre N level (+12%). Residual N levels and seasonal N release were apparently quite high, since the highest yield was with no added N plus Vitazyme (181.6 bu/acre), and yields in general tapered off as N levels increased, the lowest yields being at the 280 lb/acre N rate for both treatments. The yield response thus exhibited some N excess at the higher N rates.

Leaf pathogen results:

Northern Corn Leaf Blight Rating




Average of all plots




Vitazyme reduced the Northern Corn Leaf Blight rating significantly across nearly all plots, and did so at the 56 and 112 lb/acre N rates. At all N rates, Vitazyme treated corn reduced NCLB incidence compared to the control.

Grey Leaf Spot Rating



Average of all plots



At all levels of N except at 0 and 280 lb/acre, Vitazyme reduced the incidence of grey leaf spot. This reduction was significant at 56, 112, and 224 lb/acre of N, and also for the overall average of all plots.

Conclusions: In this North Carolina State University replicated corn study, Vitazyme increased corn grain yields at all N levels except at 56 lb/acre N. the average overall yield was increased by Vitazyme by 9%, which was significant at $P=0.05$. Resistance to both Northern Corn Leaf Blight and Grey leaf Spot were also significantly ($P=0.05$) increased over several N levels, and was significantly greater for Vitazyme over all N levels for Grey Leaf Spot, and nearly so for NCLB. This product shows excellent promise in promoting higher yields with greater disease resistance under North Carolina conditions.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2004 Crop Results

Vitazyme on Corn

Farmer: James Glass

Location: Austin, Texas

Variety: Golden Acre 2850RR

Soil type: silty clay loam

Row spacing: 30 inches

Population: 24,000 plants/acre

Planting date: March 28, 2204

Experimental design: A 110-acre field was divided into two parts, 30 acres treated with Vitazyme and the rest of the field left untreated. All other treatments were the same across the entire field.

1. Control

2. Vitazyme

Fertilization: anhydrous ammonia and a mixed N-P-K fertilizer


Vitazyme application: 13 oz/acre added to the seeds with a liquid starter fertilizer at planting

Growth observations: An examination of corn plants from each treatment at midseason revealed a noticeable size advantage for the treated plants.

Yield results: On August 27, 2004, one-acre areas of each treatment located close to each other were measured and blocked off. The one-acre areas were harvested and unloaded into a truck, and weighed individually.

Treatment	Grain weight		Change
	lb/acre	bu/acre	
Control	7,385	131.9	—
Vitazyme	7,715	137.8	5.9 (+4.5%)

Increase in grain yield: 4.5%



Conclusions: This south Texas corn study revealed that

Vitazyme applied in the seed row at planting increased the yield by 4.5% (5.9 bu/acre). There was little change in grain quality due to Vitazyme application. If a \$3.00/bushel corn price is used, this yield increase is profitable \$17.70/acre.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2004 Crop Results

Vitazyme on Corn North Carolina State University

Researcher: Ron Heiniger, Ph.D

Location: Elizabeth City, North Carolina

Variety: DKC69-71 RRH62

Tillage: conventional

Population: 33,000 seeds/acre

Row width: 30 inches

Planting date: April 21, 2004

Previous crop: soybeans

Experimental design: A plot area of 15,400 ft² (220 x 70 ft) was divided into individual plots of 400 ft² (40 x 10 ft), with four replicates. The objective of the study was to evaluate the potential of Vitazyme bios-timulant to improve grain yield and reduce disease incidence at five nitrogen rates.

Control

1. No N
2. 56 lb/acre
3. 112 lb/acre N
4. 224 lb/acre N
5. 280 lb/acre N

Vitazyme

6. No N
7. 56 lb/acre
8. 112 lb/acre N
9. 224 lb/acre N
10. 280 lb/acre N

Fertilization: 10 gal/acre of 19-19-0 %N-P₂O₅-K₂O in a 2x2 band on April 21 after broadcasting 30% UAN and a 10-34-0 fertilizer before planting

Herbicide application: Atrazine (2 qt/acre) with Banvel (0.5 pt/acre) plus Accent on May 27

Insecticide application: Counter 20CR (15 lb/acre), T-banded on April 21

Vitazyme application: 13 oz/acre on the seeds at planting, and 13 oz/acre broadcast at knee height

Harvest date: unknown

Yield Results:

Treatment	N-level	Yield	Change ¹	NCLB rating ²	Change ¹	GLS rating ³	Change ¹	
	lb/acre	bu/acre	bu/acre					
Control	0	54.0	—	3.25	—	6.13	—	
	56	118.3	—	3.75	—	6.13	—	
	112	157.8	—	3.88	—	6.38	—	
	224	150.0	—	4.25	—	6.63	—	
	280	139.9	—	4.13	—	6.75	—	
	Average		124.0	—	3.85	—	6.40	—
Vitazyme	0	98.5	44.5* (+82%)	4.88	1.63* (+50%)	6.50	0.37 (+6%)	
	56	138.4	20.4* (+17%)	5.13	1.38* (+37%)	6.88	0.75* (+12%)	
	112	155.6	(-)-2.2 (-1%)	5.13	1.25* (+32%)	7.25	0.87* (+14%)	
	224	159.4	9.4 (+6%)	5.88	1.63* (+38%)	7.38	0.75*(+11%)	
	280	161.5	21.6*(+15%)	5.75	1.62* (+39%)	7.88	1.13* (+17%)	
	Average		142.7	18.7*(+15%)	5.35	1.50* (+39%)	7.18	0.78* (+12%)
	LSD _{0.05}		15.0		0.40		0.38	


¹ Differences are compared with the same N level for control and Vitazyme

²NCLB = Northern Corn Leaf Blight: 1 = leaves covered with lesions, 10 = no disease.


³GLS = Gray Leaf Spot: 1 = leaves covered with lesions, 10 = no disease.

*Significant difference vs. the control at P=0.05.

Corn Grain Yield




Average of all plots




Vitazyme caused a substantial increase in yield over the untreated control at all N levels except at 112 lb/acre N; most of these yield differences were significant. The average yield difference was 18.7 bu/acre in favor of Vitazyme, a significant increase over the control of 15%.

Northern Corn Leaf Blight Rating




Average of all plots




At all N levels the Vitazyme treatment produced significantly reduced NCLB infection than did the control. This led to an average 39% reduction in NCLB lesions over all treatments.

Grey Leaf Spot Rating



Average of all plots



As for Northern Corn Leaf Blight ratings, Grey Leaf Spot ratings were significantly better for Vitazyme at all N levels than for the control. This differences led to a significant average difference of 12% over all plots for each treatment.

Conclusions: This corn study in North Carolina revealed that Vitazyme increased grain yield significantly, by an average of 18.7 bu/acre (+15%) over all plots ... and especially at the 0 N level, where yield was improved by 44.5 bu/acre (+82%) above the control. Both Northern Corn leaf Blight and Grey Leaf Spot were also significantly reduced by Vitazyme at all N levels, the average reduction being 39% for NCLB and 12% for GLS. These data show that Vitazyme apparently improves plant immunity to common corn pathogens, and concurrently boosts the yield potential of the crop, especially when N is limiting. The optimum N application in this study was 112 lb/acre; corn yields increased only slightly with Vitazyme with higher N rates, although without Vitazyme the yields fell somewhat.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2003 Crop Results

Vitazyme on Corn (*Surfactant vs. None*)

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Soil type: Bowie very fine sandy loam

Pot size: 1 gallon

Planting date: December 30, 2002

Experimental design: A greenhouse study was established to discover the relative effectiveness of a foliar application on corn using either diluted product in the leaf whorl or diluted product in the whorl and on leaf surfaces using a surfactant. Five replicates were used for each treatment in a complete block design. Temperatures were maintained at 55° to 80°F during the study, with no artificial light.

1. Control

2. Vitazyme in the whorl, no surfactant

3. Vitazyme on the leaves, plus a surfactant

Fertilization: All plants received 0.88 g/pot at planting of a 13-13-13% N-P₂O₅-K₂O pelleted fertilizer with 0.65% Mg, 6.0% S, 0.02% B, 0.0006% Co, 0.0006% Cu, 1.40% Fe, 0.06% Mn, 0.0006% Mo, and 0.06 % Zn. This fertilizer, giving 50 lb/acre of N, was applied to the soil surface.

Vitazyme application: On January 24, 2003, Vitazyme at 1% was sprayed from a small spray bottle into the leaf whorl of all plants in Treatment 2, being careful not to apply to the soil surface. Paper towels were used to prevent any spray from contacting the soil of the pots. Vitazyme was also sprayed the same day on the leaves and whorl of Treatment 3, with a 1% Vitazyme solution plus 5 tablespoons/gallon of Sunspray Ultra-Fine Oil, a fine agricultural oil containing 98.8% paraffinic oil.


Harvest date: February 14, 2003, 47 days after planting

Harvest results: The corn plants were washed free of soil, the leaves were measured, and then all plants were dried at about 115°F for two days. They were then weighed to the nearest 0.01 gram.

Plant Height

Treatment	Plant height*	Height change
	cm	cm
1. Control	90.3 b	—
2. Vitazyme on leaves, no surfactant	97.0 a	6.7 (+7%)
3. Vitazyme on leaves, with a surfactant	91.2 b	0.9 (+1%)


*Means followed by the same letter are not significantly different at P=0.10 according to the Tukey-Kramer Test. LSD_{0.1}=5.0 cm.



Plant Dry Weight

Treatment	Dry weight*	Weight change
	grams	grams
1. Control	7.25 b	—
2. Vitazyme on leaves, no surfactant	10.71 a	3.46 (+48%)
3. Vitazyme on leaves, with a surfactant	9.88 a	2.63 (+36%)

*Means followed by the same letter are not significantly different at $P=0.10$ according to the Tukey-Kramer Test. $LSD_{0,1}=1.58$ grams.



Conclusions: This experiment showed that corn responded almost equally well with Vitazyme applied to the leaves only, with or without a surfactant, in terms of dry weight gain during the growth period. Vitazyme in the leaf whorl only caused a highly significantly 48% weight gain versus the control, while the surfactant plus Vitazyme increased dry weight by 36%. Both treatments received the product in the leaf whorl, but Treatment 3 — with the surfactant — also had product clinging to other leaf surfaces. Both Treatments 2 and 3 had no Vitazyme applied to the soil surface.

Plant height was significantly increased by Vitazyme applied to the leaves without a surfactant, but the failure of Treatment 3 (with the surfactant) to increase significantly in height did not prevent the plants of Treatment 3 from increasing dry matter accumulation nearly as much as Treatment 2.

It is concluded from this study that, **as long as sufficient active agents are present on the plant — such as in the leaf whorl for corn — the plant will react properly to the biostimulants.** Additional amounts of product clinging to leaf surfaces as produced by a surfactant may be important in encouraging plant growth if enough droplets cling to leaf surfaces during application. However, droplets falling to the soil surface will normally contribute to product activity through root stimulation by active agents, so there may be only certain instances in which the use of a surfactant with Vitazyme may be advantageous.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2003 Crop Results

Vitazyme on Corn (*Foliar vs. Soil Application*)

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Planting rate: 10 seeds/pot thinned to 3 plants/pot

Soil type: Bowie very fine sandy loam

Planting date: December 30, 2002

Pot size: 1 gallon

Experimental design: A greenhouse study was established to discover the relative effect of soil versus foliar application of Vitazyme on corn growth. Seven replicates were set up for each treatment in a complete block design. Temperatures were maintained at 55° to 80°F during the study, with no artificial light.

1. Control

2. Vitazyme on the soil

3. Vitazyme on the leaves


Fertilization: All plants were given 0.88 g/pot at planting of a 13-13-13% N-P₂O₅-K₂O pelleted, slow release fertilizer with 0.65% Mg, 6.0% S, 0.02% B, 0.0006% Co, 0.0006% Cu, 1.40% Fe, 0.06% Mn, 0.0006% Mo, and 0.06% Zn. This fertilizer gave an effective rate of 50 lb/acre of N, applied to the soil surface.

Vitazyme application: Vitazyme was applied to the soil surface only of Treatment 2 on January 24 about at the six-leaf stage. It was also applied (a spray of a 1% solution) to the leaf whorl of the plants of Treatment 3 on January 24; care was taken to avoid applying any product to the soil surface.


Plant Height

Treatment	Plant height*	Height change
	cm	cm
1. Control	86.2 a	—
2. Vitazyme on soil	92.0 a	5.8 (+7%)
3. Vitazyme on leaves	94.5 a	8.3 (+10%)

*Means followed by the same letter are not significantly different at P=0.10 according to the Tukey-Kramer Test. LSD_{0,1}=9.7 cm.



Plant height increase (soil applied): 7%



Plant Dry Weight

Treatment	Dry weight*	Weight change
	grams	grams
1. Control	8.96 b	—
2. Vitazyme on soil	12.11 a	3.15 (+ 35%)
3. Vitazyme on leaves	10.51 ab	1.55 (+17%)

*Means followed by the same letter are not significantly different at P=0.1 according to the Tukey-Kramer Test. LSD_{0,1}=1.89 grams.

Dry weight increase (soil applied): 35%

Harvest date: February 13, 2003, 46 days after planting

Harvest results: The corn plants were washed free of soil, the leaves were measured, and then all plants were dried at about 115°F for two days, and weighed to the nearest 0.01 gram.

Conclusions: Vitazyme applied to the soil of corn in this greenhouse study produced a nonsignificant increase in plant height of 7%. Applied to the leaves, the height was increased nonsignificantly by 10%. However, Vitazyme applied to either the soil or leaves increased dry weight accumulations of the corn plants. The soil application increased growth significantly (at $P=0.10$) by 35%, and almost significantly with a foliar application (17%). It is possible that too few active agents were applied by the foliar applications for a maximum growth response, since only enough product could be applied to fill the leaf whorl; the product would not stick to the slick corn leaves. **It is concluded that both soil and foliar applications of Vitazyme are highly effective in increasing the growth rate of corn.**

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2003 Crop Results

Vitazyme on Corn

Researcher: unknown

Research organization: Department of Agriculture, Ondo State, Nigeria

Location: Iju/itaogbolu, Akure North Local Government Area, Ondo State, Nigeria

Soil type: unknown

Planting date: late season of 2000

Variety: unknown

Experimental design: A small plot replicated (3 reps), randomized complete block design was set up to evaluate the effects of Vitazyme on a number of growth parameters. Three levels of fertility were used and two applications of Vitazyme, with the following treatments:

Treatment	NPK Fertilizer	Vitazyme
1	0	yes
2	100 kg/ha	yes
3	200 kg/ha	yes
4	100 kg/ha	no
5	200 kg/ha	no
6	0	no

Fertility treatments: Treatments 2 and 4 received 100 kg/ha of an unknown fertilizer formulation two weeks after planting; Treatments 3 and 5 received 200 kg/ha of this same fertilizer also two weeks after planting.


Vitazyme treatments: Treatments 1, 2, and 3 received a 5% Vitazyme spray on the corn seeds before planting, and the newly emerged plants and soil received 1 liter/ha (13 oz/acre) two weeks after planting.

Harvest date: unknown

Growth and yield results: At harvest time several growth parameters were measured, and the data were statistically analyzed to determine significant differences at $P=0.05$.


Plant Height

Treatment	Plant height	Change
	m	m
1. (Vitazyme only)	1.5	0.3 (+25%)
2. (100 NPK + Vit.)	1.6	0.4 (+33%)
3. (200 NPK + Vit.)	1.6	0.4 (+33%)
4. (100 NPK)	1.6	0.4 (+33%)
5. (200 NPK)	1.6	0.4 (+33%)
6. (Control)	1.2	—
LSD _{0.05}	0.1	



All of the fertilizer and Vitazyme treatments significantly ($P=0.05$) increased plant height, Vitazyme alone increasing height by 25% and all other treatments increasing it by 33%.

Ears per Square Meter




Treatment	Ears number/m ²	Change number/m ²
1. (Vitazyme only)	2.7	0.4 (+17%)
2. (100 NPK + Vit.)	4.3	2.0 (+87%)
3. (200 NPK + Vit.)	4.7	2.4 (+104%)
4. (100 NPK)	4.0	1.7 (+74%)
5. (200 NPK)	4.3	2.0 (+87%)
6. (Control)	2.3	—
LSD _{0.05}	1.3	

Vitazyme alone increased ears/m² by 17%, but not significantly. However, all other Vitazyme + fertilizer treatments and all fertilizer treatments significantly increased ears/m². The Vitazyme + 200 kg/ha NPK increased ears the most, and the Vitazyme + 100 kg/ha NPK increased ears as much as did 200 kg/ha NPK, showing the ability of Vitazyme to increase the efficiency of fertilizer use.


Ear Length

Treatment	Ear length cm	Change cm
1. (Vitazyme only)	9.3	0.7 (+8%)
2. (100 NPK + Vit.)	15.3	6.7 (+78%)
3. (200 NPK + Vit.)	15.0	6.4 (+74%)
4. (100 NPK)	14.4	5.8 (+67%)
5. (200 NPK)	15.1	6.5 (+76%)
6. (Control)	8.6	—
LSD _{0.05}	1.0	



All but the Vitazyme only treatment significantly increased ear length. The Vitazyme and 100 kg/ha NPK increased ear length the most (78%), followed closely by the Vitazyme + 200 kg/ha NPK and 200 kg/ha NPK treatments.

Ear Weight




Treatment	Ear weight g	Change g
1. (Vitazyme only)	82.3	22.3 (+37%)
2. (100 NPK + Vit.)	100.7	40.7 (+68%)
3. (200 NPK + Vit.)	110.0	50.0 (+83%)
4. (100 NPK)	89.3	29.3 (+49%)
5. (200 NPK)	107.0	47.0 (+78%)
6. (Control)	60.0	—
LSD _{0.05}	11.3	

Ear weight was greatly affected by both Vitazyme alone (+37%) and by fertilizer alone (up to 78% with 200 kg/ha NPK), but most by Vitazyme + fertilizer (+68% for Vitazyme + 100 kg/ha NPK, and +83% for Vitazyme + 200 kg/ha NPK). As with ears/m² Vitazyme is shown to increase the efficiency of fertilizer use at both the 100 and 200 kg/ha NPK rates, but especially at the 100 kg/ha NPK fertilizer rate.

Grain Yield

Treatment	Grain yield g/m ²	Change* g/m ²
1. (Vitazyme only)	196.2	113.2 (+136%)
2. (100 NPK + Vit.)	338.3	255.3 (+308%)
3. (200 NPK + Vit.)	345.3	262.3 (+316%)
4. (100 NPK)	255.0	172.0 (+207%)
5. (200 NPK)	328.3	245.3 (+296%)
6. (Control)	83.0	—
LSD _{0.05}	110.3	




*All comparisons are made with the untreated control (6).

Increase in yield with Vitazyme only: 136%
Increase in yield with Vitazyme + 100 kg/ha NPK: 33%
Increase in yield with Vitazyme + 200 kg/ha NPK: 5%

All treatments significantly increased grain yield above the control. Vitazyme produced a 126% yield improvement, while the highest yield was generated by Vitazyme + 200 kg/ha NPK (+316%). This was 17.0 grams/m² higher than the 200 kg/ha NPK value. The difference was even greater for the 100 kg/ha NPK rate, where Vitazyme plus the fertilizer increased yield by 308%, but without Vitazyme the yield increased 207%. These data show a marked improvement of fertilizer efficiency with Vitazyme at the lower NPK rate, and also an improvement at the high NPK rate. These effects over the three rates are diagrammed below.

Note that the increase in grain yield above the untreated level is greatest at the lower fertilizer levels, with no fertilizer or with the 100 kg/ha NPK rate. The increase was not as dramatic at the highest NPK rate. These responses are similar to those noted in many other trials, and reflect the fact that microorganisms in the rhizosphere are stimulated to produce more available nutrients when soil nutrient levels are less than optimal. As fertility and environmental factors approach the optimum, the response from Vitazyme decreases somewhat.



Conclusions: In this replicated Nigerian corn study Vitazyme has been shown to increase plant growth and yield parameters (grain, ear number, ear length, and ear weight) significantly above the control. Vitazyme also increased yield parameters significantly, especially at the lower fertilizer levels (0 and 100 kg/ha NPK), where the Vitazyme +100 kg/ha NPK yield exceeded the 200 kg/ha NPK yield by 10.0 g/m². At 100 kg/ha NPK, Vitazyme significantly boosted yield by 83.3 g/m² above the same fertility level without Vitazyme.

In this highly weathered tropical soil of Ondo State of Nigeria, Vitazyme is seen as a powerful motivator of higher yield potential for corn.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

2003 Crop Results

Vitazyme on Corn – testimonial

Farmer/Researcher: David Schemm

Location: Arrow S Farms, Sharon Springs, Kansas

Variety: NC+ 5021RB

Planting rate: 26,000 seeds/acre

Soil type: Keith sandy clay loam

Previous crop: sunflowers

Planting date: May 3, 2003

Tillage system: no-till

Experimental design: A center pivot covering 120 acres was treated with Vitazyme over the entire area.

Fertilization: 180 lb/acre N, 35 lb/acre P₂O₅

Vitazyme and herbicide applications: (1) 13 oz/acre on May 7, with 0.5 lb/acre Atrazine 90df, 1.5 qt/acre Harness Extra, and 24 oz/acre Roundup herbicides; (2) 13 oz/acre on June 4, with 24 oz/acre Roundup herbicide when the corn was 10 inches tall

Irrigation: 16 inches total during the growing season

Weather: 8.5 inches during the growing period, with an 8-inch moisture deficit in 2002 and another 4.5 inch deficit to October of 2003; record heat throughout the summer, including several weeks of 100°F+ temperatures and 25 mph+ winds

Harvest date: October 10, 2003

Yield results: Harvested grain at 16.7% H₂O: 27,500 bushels

Yield per acre for 120 acres: 229.2 bu/acre

Conclusions: The corn received significant hail damage on June 10 when the leaves were stripped. In spite of severe heat and wind as well, the corn did exceptionally well with Vitazyme, exceeding in yield any other fields in the area. Most yields were 140 to 200 bu/acre, with a few in the 220 to 225 bu/acre range.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2002 Crop Results

Vitazyme and Awaken on Corn

Researcher: Paul W. Syltie, Ph.D.

Location: Vital Earth Resources Research Greenhouse, Gladewater, Texas

Variety: yellow dent

Soil type: Bowie very fine sandy loam

Planting date: November 21, 2001

Pot type: 1 gallon

Population: 7 seeds/pot, thinned to 3/pot

Experimental design: A complete block design was set up using eight replicates for each of four treatments. The soil was carefully packed into each pot, watered evenly, and then treated with the materials. Plants were watered on demand, and grown in the greenhouse at about 70°F for a high and 55°F for a low temperature.

1. Control

3. Awaken only

2. Vitazyme only

4. Vitazyme + Awaken

Vitazyme application: After planting on November 21, 100 ml of a 0.01% Vitazyme solution was applied to the soil surface of each pot for Treatment 2. This rate is higher than recommended, but used so as to obtain a 50-50 mixture of the two products.

Awaken application: Awaken was applied to the soil surface of the pots of Treatment 3 as 100 ml of 0.01% solution; this is equivalent to 71 oz/acre, the recommended rate for this experiment. The Awaken for Treatment 4 was mixed at the same percentage with 0.01% Vitazyme, which was also applied at 100 ml/pot.

Product specifications: Vitazyme: a liquid fermentation product of various plant materials, organisms, simple and complex carbohydrates, and other materials to yield a multiple mode of action - multiple active agent metabolic stimulator containing natural growth regulators (triacontanol, etc.), vitamins (B-complex, etc.), enzymes, and other phytoactive substances that are biologically active at very low application rates. Producer: Vital Earth Resources, Gladewater, Texas.


Awaken: a macro/micronutrient solution for plant growth stimulation having 16% N, 2% K₂O, 0.02% B, 0.15% Cu (chelated), 0.15% Fe (chelated), 0.15% Mn (chelated), 0.0006% Mo, and 2.7% Zn of which 0.15% is chelated. Awaken also contains as a major component the material called ACA (Agricultural Crop Additive). ACA's active component is zinc ammonium phosphate, the mechanism of action of which has not been fully characterized. Producer: United AgriProducts (UAP).

Harvest date: January 8, 2002, 48 days after planting.

Height results: On January 8, all of the plant roots were washed clean of soil, and the plants were measured for height. The plants were then dried in a drying oven at 115°F for 48 hours.

Treatment	Plant height*	Change vs. the control
	----- cm -----	
4. Vitazyme + Awaken	45.4 a	+11.1 (+32%)
3. Awaken	43.4 ab	+9.1 (+27%)
2. Vitazyme	40.7 b	+6.4 (+19%)
1. Control	34.3 c	—

* Means followed by the same letter are not significantly different at P=0.10, according to the Tukey-Kramer Test. LSD_{0.10}=2.0 cm.




There were some differences in plant height among the four treatments. The control treatment was significantly shorter than the other three treatments, and the Vitazyme treatment was significantly shorter than the combined Vitazyme-Awaken treatment.

Dry weight results: These results showed highly significant differences among treatment means.

Treatment	Dry weight*	Change vs. the control
	----- grams -----	
4. Vitazyme + Awaken	2.56 a	+1.20 (+30%)
3. Awaken	2.34 ab	+0.93 (+23%)
2. Vitazyme	2.09 b	+0.46 (+11%)
1. Control	1.57 c	—

* Means followed by the same letter are not significantly different at $P=0.10$, according to the Tukey-Kramer Test. $LSD_{0.10}=0.27$ g.



The dry weight of corn plants treated with Vitazyme plus Awaken was significantly greater than Vitazyme alone or the control, and also exceeded the Awaken treatment by 7%. There appears to be a synergism between Vitazyme and Awaken in this greenhouse study.

Conclusions: It appears that Vitazyme enhances the activity of Awaken for corn in terms of both plant height and dry tissue weight. Awaken, with its nutrients, stimulated corn growth somewhat more than did Vitazyme in this study, though not significantly. **Vitazyme and Awaken together appear to work well together, displaying a noticeable synergism.**

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2001 Crop Results

Vitazyme on Corn and Soybeans A Testimonial

Farmer: Tom Jones

Location: Delavan, Minnesota

Soil type: Clarion-Nicollet-Webster series

Experimental design: Several split-field experiments were set up on the Jones farm for both corn and soybeans, with variably sized treated and control areas.

Weather: Weather conditions during the year were very unfavorable for high yields, starting out very wet to delay planting, and then turning very dry for much of the summer. Yields throughout the region were down this year.

Vitazyme application: 13 oz/acre on the seeds at planting for both corn and soybeans

Data collection: Because of considerable variability in field conditions due to the wet spring and dry summer, the farmer decided not to collect yield data but closely observed effects during the season and at harvest, and during post-harvest tillage.

Chlorophyll content: Both the corn and soybeans showed more leaf chlorophyll on July 30, as detected by a Minolta SPAD chlorophyll meter. For example, one soybean trial showed the following results:

Treatment	Leaf chlorophyll*	Change
	SPAD units	
Control	35.8	—
Vitazyme	38.0	+2.2

* Twenty leaves per treatment were examined

Other crop responses noted on July 30, 2001:

Corn: taller plants, larger stalks, darker green color (more chlorophyll), larger roots with more hair roots

Soybeans: larger plants, more leaves, thicker stems, darker green color (more chlorophyll) larger roots

Observations by Tom Jones:

“Where I used Vitazyme on my beans they had more fine root hairs, and they were a little bushier in appearance. Because of the poor growing season and erratic field conditions, I didn’t get a yield check. However, they were some of my best beans.

The corn that had Vitazyme on also had a lot more fine root hairs. I couldn’t believe the difference when I disced my stalks. I could see all these bushy looking root balls, unlike in the untreated fields. Again, no yield check was taken, but I know I could see a difference in the combine hopper. I plan to use more Vitazyme next year – perhaps on all my acres.”

Thanks,

Tom Jones

Faribault County

Southern Minnesota

Effects of Solution pH During Storage On Vitazyme Efficacy With Corn (*Zea Mays L.*)

*By Paul W. Syltie, Ph.D., Soil Fertility
Director of Research, Vital Earth Resources
706 East Broadway Avenue, Gladewater, Texas 75647, U.S.A.
June, 2001*

Introduction

Vitazyme is a naturally fermented biostimulant that contains a multiple array of active agents — vitamins, enzymes, growth regulators, and other substances — which trigger various growth responses in plants. Typical effects include enhanced root growth, greater leaf chlorophyll, increased carbon fixation, and concomitant increases in overall growth, life-cycle stimulation, root exudation, rhizosphere microbial growth, and crop yields.

Little information is available regarding the effects of the pH of the solution during storage on the resultant effectiveness of Vitazyme's active agents for plant growth. Therefore, this study, using corn as the test crop, was initiated to answer the questions of efficacy with pH over time.

Materials and Methods

Two Vitazyme concentrations were used in this study — 1% and 100% — to simulate conditions during use in the field when mixed either undiluted or diluted with agricultural chemicals. These solutions were placed in beakers which were sealed with Parafilm to prevent evaporation. For each concentration, the pH of the solution was adjusted to pH 7.0, 8.0, 9.0, or 10.0. These dilutions, prepared on April 4, 2001, are summarized in Table 1. The pH of each solution was determined again on May 7, 2001, 34 days after initial preparation.

Table 1. Dilutions of Vitazyme at various pH levels for a corn efficacy study.

Solution pH*	Parts of Vitazyme : Parts of Water (distilled)	
	100% solution	1% solution
7.0	100:0	1:99
8.0	100:0	1:99
9.0	100:0	1:99
10.0	100:0	1:99

* Solution pH was adjusted using a NaOH solution having a pH of 12.90

On May 8, 2001, the corn study was initiated in the Vital Earth Resources research greenhouse. One gallon pots were filled with Bowie fine sandy loam and placed in a complete block arrangement (eight replications), with five treatments for each concentration. See Table 2 for a summary of these treatments.

Table 2. Treatments for corn in a Vitazyme study using two concentrations of product stored at various pH's.

Treatment	Vitazyme, 100%	Vitazyme, 1%
1. Control (no Vitazyme)	none	none
2. pH 7.0	x	x
3. pH 8.0	x	x
4. pH 9.0	x	x
5. pH 10.0	x	x

Seven corn seeds (yellow dent, treated with Captan fungicide) were planted in each pot at a depth of 0.75 inch, and each pot received 100 ml of solution carefully distributed to the soil surface of the pots. The 100% Vitazyme pots received 100 ml of a 0.1% (1 ml/liter) solution of actual Vitazyme, while the 1% Vitazyme pots received 100 ml of a 0.05% (50 ml/liter) solution of actual Vitazyme. The 1% solution was applied half as concentrated as the 100% solution because there was not enough prepared solution of the 1% concentration.

On May 5, 2001, the emerged corn plants were thinned to three aggressive plants per pot, and on May 30, 2001, 22 days after planting, the plants were harvested. All soil was washed from the roots, the height of each plant was measured, and the plants were dried in a drying oven at about 115° F for two days. Each set of three plants from each pot was weighed to the nearest 0.01 gram, and a statistical analysis (ANOVA) was run on each concentration (100% and 1%) using Cohort software.

Results and Discussion

Solution pH changes over 34 days

The pH of the stored solutions tended to move towards neutrality over the 34-day storage period. Interestingly, the 1% concentration moved more towards neutrality than did the 100% concentration (see Figure 1).




Figure 1. pH changes during 34 days of storage of Vitazyme solutions.

Vitazyme tended to move towards neutrality (pH 7.0) at both the 1% and 100% concentrations when stored at room temperature for 34 days. This was equally true for the dilute (1%) solution, where all four solution pH's ended up between pH 7.08 and 7.66 at the end of the storage period.

Corn Growth Effects

The average height and weight of the corn plants for the treatments are shown in Table 3.

Table 3. Corn heights and weights treated with different Vitazyme solutions stored at various pH's.

Treatment	Corn height (cm)	Tissue dry weight (g)
1% Vitazyme		
1. Control	39.00	1.95
2. pH 7.0	44.88	2.64
3. pH 8.0	46.13	2.81
4. pH 9.0	45.00	2.73
5. pH 10.0	41.88	2.37
100% Vitazyme		
1. Control	45.38	2.59
2. pH 7.0	48.00	2.79
3. pH 8.0	48.25	2.97
4. pH 9.0	47.75	2.71
5. pH 10.0	45.63	2.70

A statistical analysis of the data revealed the following, as shown in Table 4.

Table 4. A statistical analysis of a corn study, using Vitazyme stored at different pH's.

Treatment	Corn height (cm)	Tissue dry weight (g)
1% Vitazyme*		
3. pH 8.0	46.13 a (+18%)	2.81 a (+44%)
4. pH 9.0	45.00 a (+15%)	2.73 a (+40%)
2. pH 7.0	44.88 ab (+15%)	2.64 ab (+35%)
5. pH 10.0	41.88 bc (+7%)	2.37 b (+22%)
1. Control	39.00 c	1.95 c
CV	5.48%	10.70%
LSD _{0.10}	2.02	0.23
100% Vitazyme*		
3. pH 8.0	48.25 a (+6%)	2.97 a (+15%)
2. pH 7.0	48.00 a (+6%)	2.79 ab (+8%)
4. pH 9.0	47.75 a (+5%)	2.71 ab (+5%)
5. pH 10.0	45.63 a (+1%)	2.70 ab (+4%)
1. Control	45.38 a	2.59 b
CV	4.78%	9.51%
LSD _{0.10}	1.91	0.22

*Means followed by the same letter are not significantly different according to the Tukey-Kramer Test (P=0.10).

It is clear that Vitazyme in every case, at both the 1% and 100% concentrations, and at all pH's, provided height and dry weight increases in this corn study. Several of these increases were significant at $P=0.01$, especially when Vitazyme was stored at pH 8.0 for both the 1% and 100% concentrations. At 1%, the pH 8.0 solution provided a 44% dry weight increase for Vitazyme, and at 100% a 15% increase. Plant height increases were in all cases less than half of the dry weight increases, but followed the same order as with dry tissue weight.

Vitazyme stored at pH 7.0, 8.0, and 9.0 always provided good dry weight increases for corn, especially for the 1% solution. At pH 10.0, however, the increases were less — indicating some deactivation of active agents — though with the 1% solution the increase was still significant; for the 100% solution the increase was not significantly greater than with the untreated control.

Summary and Conclusions

Vitazyme at 1% dilution, when adjusted to pH 7.0, 8.0, or 9.0, and stored for 34 days at room temperature, always caused significant height and weight increases for corn in this study. Increases in dry weight were up to 44% above the control. At pH 10.0 the increases were smaller but, in the case of dry weight, still significantly greater than the control. At the 100% dilution the increase in dry weight with Vitazyme at pH 8.0 was significant, and all other pH's also produced increases.

This study reveals that Vitazyme significantly improved corn height and dry weight at any product pH when stored for 34 days, but especially at pH 8.0. Thus, the use of Vitazyme in fertilizer or pesticide solutions of pH 7.0 to 9.0 is recommended, and such use can boost product efficacy. Product effectiveness when stored at pH 10.0 appears to be somewhat diminished.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2001 Crop Results

Vitazyme on Corn

Researcher: Dennis Parrett, Cecilia Farm Service, Inc.

Location: Cecelia, Kentucky

Farm cooperater: Richard Preston

Variety: Novartis 6367

Row spacing: 30 in

Population: 26,600 seeds/acre

Planting date: April 10, 2001

Soil type: unknown

Experimental design: A test field was divided into five portions, each with a treatment as shown below.

Treatment	Foliar N	Sidedress N	Vitazyme
1. Control	0	0	0
2. Foliar N	5 gal/acre of 28% N	0	0
3. Sidedress N, low	0	80 lb/acre of 28% N	0
4. Sidedress N, high	0	105 lb/acre of 28% N	0
5. Sidedress N, high + Vitazyme	0	105 lb/acre of 28% N	13 oz/acre



Fertilization: 50-60-60 lb/acre N-P₂O₅-K₂O preplant incorporated; sidedress fertilizer as shown above

Vitazyme treatments: 13 oz/acre along with the 28% nitrogen solution, applied June 9

Yield and income results: Harvest was on October 11, 2001.

Treatment	Grain yield	Increase over the control	Product cost*	Net return
	bu/acre	bu/acre	\$/acre	\$/acre
1. Control	102.6	—	0	—
2. Foliar N	112.3	9.7 (+9%)	15.88	5.46
3. Sidedress N, low	124.4	21.8 (+21%)	33.25	44.85
4. Sidedress N, high	138.1	35.5 (+35%)	26.52	21.48
5. Sidedress N, high + Vitazyme	151.9	49.3 (+48%)	37.25	61.27

* Product costs were determined by Cecilia Farm Service, including \$4.00/acre for Vitazyme and \$2.00/bu for corn.



True increase from Vitazyme

Yield: Treatment 5 vs. Treatment 4: 151.9 bu/acre – 138/bu/acre = **13.8 bu/acre**

Net income: Treatment 5 vs Treatment 4: \$61.27/acre – \$21.48/acre = **\$39.79/acre**

Conclusions: In this corn study in Kentucky, Vitazyme proved to be highly beneficial to corn production in terms of yield increase and income increase. The 13.8 bu/acre increase in yield provided an extra \$39.79/acre income, showing its high profitability in farming programs.

Return per dollar invested in Vitazyme: \$9.95

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2001 Crop Results

Vitazyme on Sweet Corn

New York Crop Research Facility, Cornell University

Researchers: Arlie McFaul, Alan Erb, Lee Stivers, and Christy Hoeping

Location: near Batavia, New York

Variety: Bonus

Row spacing: 30 inches

Spacing-in-row: 9 inches

Planting date: June 4, 2001

Experimental design: A small field experiment was designed in a randomized complete block fashion, with four replications. Individual plots were 6 rows wide and 20 feet long. Foliar treatments were made to the center two rows only for all eight treatments.

1. Vitazyme
2. Harpin protein seed treatment
3. Messenger
4. ACA

5. Asset RS
6. Auxigrow
7. K-Mag
8. Control

Fertilization: All areas received 250 lb/acre of a 15-15-15% N-P₂O₅-K₂O dry formulation banded along the seed row at planting. On July 10, 100 lb/acre of N was applied.

Vitazyme application: (1) 13 oz/acre sprayed over the soil after emergence on July 6, and (2) again before tasseling on July 30; other products were added according to supplier recommendations.


Herbicides: Atrazine (3 pints/acre) and Basagran (1.5 pints/acre) post-emergent on June 21

Harvest date: August 31, 2001

Weather: It was very hot and dry during the summer, with growth and yields curtailed due to the drought. Monthly totals: May, 3.84 in; June, 1.47 in; July, 1.02 in; August 2.21 in; September, 2.82 in.

Yield and harvest results: Total yield and cob characteristics were evaluated before or at harvest



Plant population



Percent Tip Fill

All values were not significantly different, but the population with Vitazyme was the highest, being 7% higher than the control. Some significant differences in percent tip fill appeared, with Messenger and Asset RS being the lowest in value of all treatments. Vitazyme was statistically equal to the highest tip fill value.

*Means followed by the same letter are not significantly different according to Fisher's Protected LSD; $P_{0.05} = 0.04$.





Ear Diameter

There was relatively little difference amongst the various treatments for ear diameter. None were significantly different.

Number of Rows

Vitazyme and K-Mag had the highest numbers of rows of kernels per ear, being 2% higher than the control. There were no significant differences amongst these values.





Ears per Acre

Vitazyme produced the greatest number of ears per acre, being slightly greater than the control but considerably greater than the other treatments by about 13%. This increase was a reflection of higher estimated plant population as noted earlier. No treatment means were significantly different.

Corn Yield

Vitazyme yielded the most corn of any treatment, exceeding the control by 5% but exceeding the lowest other treatment (ACA) by 33%. The difference was equivalent to \$54.88/acre based on a \$49/ton sweet corn price. None of these differences were significant, however.



Cob Weight, 35 Ears

* Growers are urged to grow large ears, so the weight of 35 ears should equal or exceed 25 lb; lower prices result from underweight ears. In 2001 the weights were low because of the severe drought. Nonetheless, the 25-ear weight for Vitazyme was the highest of all the treatments, exceeding the control by 4%. Vitazyme exceeded ACA ear weight by 17%.

Conclusions: Results with sweet corn in this study were greatly affected by a severe summer drought. In spite of this fact, **Vitazyme performed the best of all seven treatments used in this study, being highest in plant population, rows per ear, ears per acre, yields per acre, and total weight per 35 ears. Vitazyme treatment produced 33% more yield than the lowest yielding other treatment.** Though significance in the mean differences was lacking, the consistent trend of this study was for Vitazyme to provide excellent plant responses that would substantially benefit sweet corn growers.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

2000 Crop Results

Vitazyme on Corn

Grower: Jim Clise

Location: Waterloo, New York

Variety: Agway 501

Researcher: Jake Gephart, Agway, Inc.

Planting date: May 31, 2000

Seeding rate: unknown

Row spacing: 30 inches

Experimental design: A 100-acre field was divided into halves (50 acres each), with half treated with Vitazyme and half left untreated.

1. Control

2. Vitazyme

Fertilization: 380 lb/acre of a urea and potash mixture, broadcast before planting and incorporated; 250 lb/acre of 11-37-0% N-P₂O₅-K₂O as a starter

Weed control: Bicep and Prowl tank-mixed


Vitazyme application: 13 oz/acre with the herbicide, at 18-inches corn height

Harvest date: December 3, 2000

Yield results:

	Control	Vitazyme	Change
	bu/acre		
Corn yield	141	155	(+) 14 (+10%)

Yield increase: 10%



Income results:

	Control	Vitazyme	Change
	\$/acre		
Income	246.75	271.25	(+) 24.50

**Income increase:
\$24.50/acre**

Conclusions: Only 13 oz/acre of Vitazyme, applied with the herbicide, resulted in a 10% grain yield increase and a \$24.50/acre income increase. This increase resulted in a return on investment of about 5:1 for this low-value crop. In spite of a very wet and cool year, Vitazyme still produced a very good crop response.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

1999 Crop Results

Vitazyme on Corn

Farmer: Craig Rice, Rice Farms

Location: Geneseo, New York

Supervisor: Don Jones, Agway Inc.

Variety: unknown

Planting date: unknown

Harvest date: November 10, 1999

Experimental design: A large field was divided in two, with part treated with Vitazyme and part left untreated. Yields were determined by harvesting a 15- ft wide strip that was 254 to 768 feet long for each treatment.

1. Control


2. Vitazyme

Fertility treatments: unknown

Vitazyme treatment: 13 oz/acre at planting

Chlorophyll determinations: On August 12 readings were taken of leaves from the corn treatments with a Minolta SPAD chlorophyll meter. Each value represents an average of 10 individual leaf determinations.


Treatment	SPAD units	Increase
Control	54.7	--
Vitazyme	56.7	2.0



**Chlorophyll increase:
2.0 SPAD units**

Yield results:

Treatment	Yield lb/plot	Plot area acre	Yield bu/plot
Control	2,700	0.264	182.6
Vitazyme, area 1	2,865	0.259	197.5
Vitazyme, area 2	2,690	0.259	185.4
Vitazyme, average	---	---	191.5



Yield increase: 5%

Income increase: Estimated corn price = \$2.50/bu.

8.9 bu/acre increase x \$2.50/bu = \$22.25/acre

Income increase: \$22.50/acre

Comments: In spite of a very dry and hot summer, the Vitazyme treatment boosted the corn yield significantly (8.9 bu/acre). Leaf chlorophyll increases during the growing season would explain most of this increase, since this would promote greater root growth and exudation to feed a more vigorous rhizosphere organism population.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

1999 Crop Results

Vitazyme on Corn

Farmer: Bill Goodell

Location: Shortsville, New York

Variety: Pioneer 3573

Planting date: May 4, 1999

Population: 30,000 seeds/acre planted, 28,000 plants/acre final

Experimental design: A 5-acre field was split in half, one half receiving Vitazyme with 50% of the usual starter fertilizer at planting, and the other half receiving the regular starter rate but no Vitazyme.

1. Control: 100% starter

2. Vitazyme + 50% starter

Fertility treatments: All areas of the field received a broadcast application of 100 lb/acre $(\text{NH}_4)_2\text{SO}_4$ (21%N) + Boron + Copper, 500 lb/acre EnviroSoil (composted sewage sludge mixed with high-calcium lime), 32 gal/acre 30% UAN, and 1 gal/acre liquid Ca-nitrate. At planting the control received 10 gal/acre of 3-18-18 (\$3.25/gal = \$32.50/acre), while the Vitazyme treated area received 5 gal/acre of 3-18-18 (\$16.25/acre). Total N per acre for the broadcast fertilizer was 107 lb/acre, excluding the compost, for the control, and 106 lb/acre for the Vitazyme treatment.


Vitazyme treatment: 13 oz/acre mixed with the 3-18-18, on the seeds at planting

Weather conditions: very dry season-long

Yield results:

<u>Treatment</u>	<u>Yield</u> bu/acre	<u>Test weight</u> lb/bu
Control	76.12	57.0
Vitazyme	78.78 (+3.5%)	57.5

Test weight increase: 0.5 lb/bu



Yield increase with 50% starter: 3.5 %

Income results: Corn is priced at \$2.00/bu.

<u>Treatment</u>	<u>Corn value</u>	<u>Fertilizer savings*</u>	<u>Total Increase</u>
Control	\$152.24/acre	---	---
Vitazyme	\$157.56/acre	\$12.35/acre	\$17.67/acre

Comments: Yields were reduced by about 50%, due to very dry conditions throughout the summer. In spite of this, Vitazyme stimulated a yield increase with a reduced starter fertilizer input, illustrating its ability to activate rhizosphere nutrient uptake.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647

(903) 845-2163 FAX: (903) 845-2262

1997 Crop Results

Vitazyme on Corn

Researcher: William (Bill) Goodell

Seeding date: May 14, 1997

Location: Shortsville, New York

Row width: 30 inches

Seeding rate: 30,000 seeds/acre

Variety: Pioneer 3752 (97 day)

Experimental design: A field of reasonable uniformity was divided into two parts: an untreated control part and a Vitazyme treated part.

1. Control (no Vitazyme)

2. Vitazyme + 33% of fertilizer at planting

Fertility treatments: The entire field received 10 tons/acre of manure, plus 26 gal/acre of 30% nitrogen and 4 gal/acre of ammonium thiosulfate. The control area received an additional 8 gal/acre of 9-18-9 plus 4 gal/acre of 0-0-30 at planting on the seed. The Vitazyme treatment received 2.7 gal/acre of 9-18-9 plus 1.3 gal/acre of 0-0-30 at planting, or 33% of the control treatment.

Vitazyme application: 12 oz/acre with the liquid fertilizer, on the seeds at planting

Soil: Ontario loam, 3 to 10% slope

Previous crop: wheat

Harvest date: November 26, 1997

Yield results: Both treatments yielded about 22.7% grain moisture.

	<u>Control</u>	<u>Vitazyme</u>	<u>Increase with Vitazyme</u>
Corn yield	129.6 bu/acre	142.2 bu/acre	12.6 bu/acre

**Yield Increase:
10%**

Income results: The grain price has been calculated at about \$3.00/bu. Natures 9-18-9 + 0-0-30, mixed at a 2:1 ratio retails for about \$3.20/gal.

	<u>Control</u>	<u>Vitazyme</u>	<u>Income Increase</u>
Corn income	\$386.12/acre	\$423.64/acre	\$37.80/acre
Fertilizer savings, less Vitazyme	—	\$25.60/acre	\$25.60/acre
Net income increase			\$63.40/acre

Income Increase: \$63.40/acre

Test weight results:

	<u>Control</u>	<u>Vitazyme</u>
Test weight	53 lb/bu	55 lb/bu

**Test weight
increase: 2 lb/bu**

Comments: The cropping year was quite good. Only July 13, leaf chlorophyll measured on 20 average leaves of each treatment gave 50.7 SPAD units for the Vitazyme treatment and 49.1 SPAD units for the control.

Vital Earth Resources

706 East Broadway, Gladewater, Texas 75647
(903) 845-2163 FAX: (903) 845-2262

1997 Crop Results

Vitazyme on Corn

Researcher: William (Bill) Goodell

Seeding date: May 14, 1997

Location: Shortsville, New York

Row width: 30 inches

Seeding rate: 30,000 seeds/acre

Variety: Pioneer 3752 (97 day)

Experimental design: A field of reasonable uniformity was divided into two parts: an untreated control part and a Vitazyme treated part.

1. Control (no Vitazyme)

2. Vitazyme + 33% of fertilizer at planting

Fertility treatments: The entire field received 10 tons/acre of manure, plus 26 gal/acre of 30% nitrogen and 4 gal/acre of ammonium thiosulfate. The control area received an additional 8 gal/acre of 9-18-9 plus 4 gal/acre of 0-0-30 at planting on the seed. The Vitazyme treatment received 2.7 gal/acre of 9-18-9 plus 1.3 gal/acre of 0-0-30 at planting, or 33% of the control treatment.

Vitazyme application: 12 oz/acre with the liquid fertilizer, on the seeds at planting

Soil: Ontario loam, 3 to 10% slope

Previous crop: wheat

Harvest date: November 26, 1997

Yield results: Both treatments yielded about 22.7% grain moisture.

	<u>Control</u>	<u>Vitazyme</u>	<u>Increase with Vitazyme</u>
Corn yield	129.6 bu/acre	142.2 bu/acre	12.6 bu/acre

**Yield Increase:
10%**

Income results: The grain price has been calculated at about \$3.00/bu. Natures 9-18-9 + 0-0-30, mixed at a 2:1 ratio retails for about \$3.20/gal.

	<u>Control</u>	<u>Vitazyme</u>	<u>Income Increase</u>
Corn income	\$386.12/acre	\$423.64/acre	\$37.80/acre
Fertilizer savings, less Vitazyme	—	\$25.60/acre	<u>\$25.60/acre</u>
Net income increase			\$63.40/acre

Income Increase: \$63.40/acre

Test weight results:

	<u>Control</u>	<u>Vitazyme</u>
Test weight	53 lb/bu	55 lb/bu

**Test weight
increase: 2 lb/bu**

Comments: The cropping year was quite good. Only July 13, leaf chlorophyll measured on 20 average leaves of each treatment gave 50.7 SPAD units for the Vitazyme treatment and 49.1 SPAD units for the control.